

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
16 June 2005 (16.06.2005)

PCT

(10) International Publication Number
WO 2005/053516 A2

- (51) International Patent Classification⁷: **A61B**
- (21) International Application Number:
PCT/US2004/040120
- (22) International Filing Date: 1 December 2004 (01.12.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/525,989 1 December 2003 (01.12.2003) US
- (71) Applicant (*for all designated States except US*): **BOARDS OF REGENTS, THE UNIVERSITY OF TEXAS SYSTEM** [US/US]; 201 West 7th Street, Austin, TX 78701 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **LEMON, Stanley, M.** [US/US]; 1517 Bayou Shore Drive, Galveston, TX 77551 (US). **YI, MinKyung** [KR/US]; 7700 Seawall Blvd.#301, Galveston, TX 77551 (US).
- (74) Agent: **PROVENCE, David, L.**; Mueting, Raasch & Gebhardt, P.A., P.O. Box 581415, Minneapolis, MN 55454-1415 (US).
- (81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: REPLICATION COMPETENT HEPATITIS C VIRUS AND METHODS OF USE

(57) Abstract: The present invention provides replication competent polynucleotides that include a coding sequence encoding a hepatitis C virus polypeptide having adaptive mutations. The invention also includes methods for making replication competent polynucleotides, identifying a compound that inhibits replication of a replication competent polynucleotide, selecting a replication competent polynucleotide, and detecting a replication competent polynucleotide.



WO 2005/053516 A2

5 REPLICATION COMPETENT HEPATITIS C VIRUS
 AND METHODS OF USE

 CONTINUING APPLICATION DATA

 This application claims the benefit of U.S. Provisional Application Serial
10 No. 60/525,989, filed December 1, 2003, which is incorporated by reference
 herein.

 GOVERNMENT FUNDING

 The present invention was made with government support under Grant
15 Nos. U19-AI40035 and N01-AI25488, awarded by the National Institute of
 Allergy and Infectious Diseases. The Government has certain rights in this
 invention.

 BACKGROUND

20 Hepatitis C virus is the most common cause of chronic viral hepatitis
 within the United States, infecting approximately 4 million Americans and
 responsible for the deaths of 8,000-10,000 persons annually due to progressive
 hepatic fibrosis leading to cirrhosis and/or the development of hepatocellular
 carcinoma. Hepatitis C virus is a single stranded, positive-sense RNA virus with a
25 genome length of approximately 9.6 kb. It is currently classified within a separate
 genus of the flavivirus family, the genus *Hepacivirus*. The hepatitis C virus
 genome contains a single large open reading frame (ORF) that follows a 5' non-
 translated RNA of approximately 342 bases containing an internal ribosome entry
 segment (IRES) directing cap-independent initiation of viral translation. The
30 large ORF encodes a polyprotein which undergoes post-translational cleavage,
 under control of cellular and viral proteinases. This yields a series of structural
 proteins which include a core or nucleocapsid protein, two envelope
 glycoproteins, E1 and E2, and at least six nonstructural replicative proteins.
 These include NS2 (which with the adjacent NS3 sequence demonstrates *cis*-
35 active metalloproteinase activity at the NS2/NS3 cleavage site), NS3 (a serine
 proteinase/NTPase/RNA helicase), NS4A (serine proteinase accessory factor),
 NS4B, NS5A, and NS5B (RNA-dependent RNA polymerase).

With the exception of the 5' non-translated RNA, there is substantial genetic heterogeneity among different stains of hepatitis C virus. Phylogenetic analyses have led to the classification of hepatitis C virus strains into a series of genetically distinct "genotypes," each of which contains a group of genetically related viruses. The genetic distance between some of these genotypes is large enough to suggest that there may be biologically significant serotypic differences as well. There is little understanding of the extent to which infection with a virus of any one genotype might confer protection against viruses of a different genotype.

10 The currently available therapy of interferon in combination with ribavirin has poor response rate against most prevalent strains of HCV, genotype 1a and 1b. Establishment of selectable subgenomic replicon systems has advanced the study of HCV RNA replication. However, only replicons of genotype 1b strains are readily available, and extension of replicon systems to other genotypes has been largely
15 unsuccessful. Considering the nature of high genetic variability of HCV, HCV replication systems derived from other genotypes will be very helpful in the effort of drug discovery. In support with this notion, chimeric replicons containing a genotype 1a polymerase in the background of a genotype 1b replicon were more resistant to interferon treatment in vitro than the replicon derived from a genotype 1b HCV.
20 Extension of replicon system to other genotypes are also necessary to understand the mechanism of HCV RNA replication and the contribution of variable sequences in that process.

 Recently two groups reported the generation of genotype 1a replication system using highly permissive sublines of Huh-7 cells. Blight et al. (J. Virol. 77,
25 3181-3190 (2003)) were able to select G418 resistant colonies supporting replication of genotype 1a derived subgenomic replicons in a hyper-permissive Huh7 subline, Huh-7.5, that was generated by curing an established G418-resistant replicon cell line of the subgenomic Con1 replicon RNA that had been used to select it by treatment with interferon-alpha (Blight et al., J. Virol., 76,
30 13001-13014 (2002)). Sequence analysis of replicating HCV RNAs inside of such selected cell lines showed that the most common critical mutations were located at amino acid position 470 of NS3 (P1496L) within domain II of the NS3 helicase, and the NS5A mutation (S2204I). In other case, Grobler et al. (J. Biol. Chem., 278,16741-16746 (Feb, 2003)), used a systematic mutational approach to reach

the similar conclusion that both P1496L and S2204I combination was necessary to get genotype 1a replication in a highly permissive Huh-7 subline which was selected in an independent but similar way. However, genotype 1a RNAs with these two enhanced mutations does not undergo replication in the Huh-7 cell line, indicating limited usefulness of this system.

SUMMARY

The present invention provides replication competent polynucleotides. The replication competent polynucleotides include a 5' non-translated region (NTR), a 3' NTR, and a first coding sequence present between the 5' NTR and 3' NTR and encoding a hepatitis C virus polyprotein. The 5' NTR, the 3' NTR, and the nucleotide sequence encoding the polyprotein may be genotype 1a. The polyprotein includes an isoleucine at about amino acid 2204, and further includes an adaptive mutation. The adaptive mutation can be an arginine at about amino acid 1067, an arginine at about amino acid 1691, a valine at about amino acid 2080, an isoleucine at about amino acid 1655, an arginine at about amino acid 2040, an arginine at about amino acid 1188, or a combination thereof. The polyprotein may be a subgenomic polyprotein. The polyprotein may include the cleavage products core, E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B.

The replication competent polynucleotides may further include a second coding sequence. The second coding sequence can encode, for instance, a marker or a transactivator. The replication competent polynucleotides may further include a nucleotide sequence having cis-acting ribozyme activity, wherein the nucleotide sequence is located 3' of the 3' NTR.

Also provided by the present invention are methods for making a replication competent polynucleotide, and the resulting replication competent polynucleotide. The methods include providing a polynucleotide having a 5' NTR, 3' NTR, a first coding sequence present between the 5' NTR and 3' NTR and encoding a hepatitis C virus polyprotein. Typically, the 5' NTR, polyprotein, and 3' NTR are genotype 1a. The polyprotein includes a serine at about amino acid 2204, a glutamine at about amino acid 1067, a lysine at about amino acid 1691, a phenylalanine at about amino acid 2080, a valine at about amino acid 1655, a lysine at about amino acid 2040, or a glycine at about amino acid 1188. The

method also includes altering the coding sequence such that the polyprotein encoded thereby includes an isoleucine at amino acid 2204, and an adaptive mutation. The polyprotein may be a subgenomic polyprotein. The polyprotein may include the cleavage products core, E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B.

The present invention further provides methods for identifying a compound that inhibits replication of a replication competent polynucleotide. The method includes contacting a cell containing a replication competent polynucleotide with a compound, incubating the cell under conditions wherein the replication competent polynucleotide replicates in the absence of the compound, and detecting the replication competent polynucleotide, wherein a decrease of the replication competent HCV polynucleotide in the cell contacted with the compound compared to the replication competent polynucleotide in a cell not contacted with the compound indicates the compound inhibits replication of the replication competent polynucleotide. The detecting of the replication competent polynucleotide can include, for instance, nucleic acid amplification or identifying a marker encoded by the replication competent polynucleotide or by the cell containing the replication competent polynucleotide.

Also provided by the present invention are methods for selecting a replication competent polynucleotide. The method includes incubating a cell containing a polynucleotide including a 5' NTR, a 3' NTR, and a first coding sequence present between the 5' NTR and 3' NTR and encoding a hepatitis C virus polyprotein, and a second coding sequence. The polyprotein includes an isoleucine at about amino acid 2204, and further includes an adaptive mutation. The second coding sequence encodes a selectable marker conferring resistance to a selecting agent that inhibits replication of a cell that does not express the selectable marker. The method also includes detecting a cell that replicates in the presence of the selecting agent, wherein the presence of such a cell indicates the polynucleotide is replication competent. The method may further include obtaining a virus particle produced by the cell, exposing a second cell to the isolated virus particle and incubating the second cell in the presence of the selecting agent, and detecting a second cell that replicates in the presence of the selecting agent, wherein the presence of such a cell indicates the replication competent polynucleotide in the first cell produces an infectious virus particle.

The present invention also provides methods for detecting a replication competent polynucleotide, including incubating a cell containing a replication competent polynucleotide. The replication competent polynucleotide includes a 5' NTR, a 3' NTR, and a first coding sequence present between the 5' NTR and 3' NTR and encoding a hepatitis C virus polyprotein, and a second coding sequence encoding a transactivator. The cell includes a transactivated coding region and an operator sequence operably linked to the transactivated coding region, and the transactivated coding region encodes a detectable marker, wherein the transactivator alters transcription of the transactivated coding region. The method further includes detecting the detectable marker, wherein the presence of the detectable marker indicates the cell includes a replication competent polynucleotide.

Definitions

As used herein, the term "replication competent polynucleotide" refers to a polynucleotide that replicates when present in a cell. For instance, a complementary polynucleotide is synthesized. As used herein, the term "replicates *in vitro*" indicates the polynucleotide replicates in a cell that is growing in culture. The cultured cell can be one that has been selected to grow in culture, including, for instance, an immortalized or a transformed cell. Alternatively, the cultured cell can be one that has been explanted from an animal. "Replicates *in vivo*" indicates the polynucleotide replicates in a cell within the body of an animal, for instance a primate (including a chimpanzee) or a human. In some aspects of the present invention, replication in a cell can include the production of infectious viral particles, i.e., viral particles that can infect a cell and result in the production of more infectious viral particles.

As used herein, the term "polynucleotide" refers to a polymeric form of nucleotides of any length, either ribonucleotides or deoxynucleotides, and includes both double- and single-stranded DNA and RNA. A polynucleotide may include nucleotide sequences having different functions, including for instance coding sequences, and non-coding sequences such as regulatory sequences and/or non-translated regions. A polynucleotide can be obtained directly from a natural source, or can be prepared with the aid of recombinant, enzymatic, or chemical techniques. A polynucleotide can be linear or circular in topology and can be, for

example, a portion of a vector, such as an expression or cloning vector, or a fragment.

The terms "coding region" and "coding sequence" are used interchangeably and refer to a polynucleotide region that encodes a polypeptide and, when placed under the control of appropriate regulatory sequences, expresses the encoded polypeptide. The boundaries of a coding region are generally determined by a translation start codon at its 5' end and a translation stop codon at its 3' end. A coding region can encode one or more polypeptides. For instance, a coding region can encode a polypeptide that is subsequently processed into two or more polypeptides. A regulatory sequence or regulatory region is a nucleotide sequence that regulates expression of a coding region to which it is operably linked. Nonlimiting examples of regulatory sequences include promoters, transcription initiation sites, translation start sites, internal ribosome entry sites, translation stop sites, and terminators. "Operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A regulatory sequence is "operably linked" to a coding region when it is joined in such a way that expression of the coding region is achieved under conditions compatible with the regulatory sequence.

"Polypeptide" as used herein refers to a polymer of amino acids and does not refer to a specific length of a polymer of amino acids. Thus, for example, the terms peptide, oligopeptide, protein, polyprotein, proteinase, and enzyme are included within the definition of polypeptide. This term also includes post-expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like. A "hepatitis C virus polyprotein" refers to a polypeptide that is post-translationally cleaved to yield more than one polypeptide.

The terms "5' non-translated RNA," "5' non-translated region," "5' untranslated region" and "5' noncoding region" are used interchangeably, and are terms of art (see Bukh et al., Proc. Nat. Acad. Sci. U S A, 89, 4942-4946 (1992)). The term refers to the nucleotides that are at the 5' end of a replication competent polynucleotide.

The terms "3' non-translated RNA," "3' non-translated region," and "3' untranslated region" are used interchangeably, and are terms of art. The term

refers to the nucleotides that are at the 3' end of a replication competent polynucleotide.

Unless otherwise specified, "a," "an," "the," and "at least one" are used interchangeably and mean one or more than one.

5

BRIEF DESCRIPTION OF THE FIGURES

Figure. 1. Organization of the selectable subgenomic dicistronic HCV replicons, Bpp-Ntat2ANeo/SI (identical to Ntat2ANeo/SI in Yi et al., Virol., 302, 197-210 (2002)), Htat2ANeo/SI, and Bpp-Htat2ANeo/SI, in which most of the nonstructural protein-coding region and the 3'NTR are derived from the H77c HCV genotype 1a sequence. The two large ORFs are shown as rectangles, with nontranslated RNA segments shown as lines. The segment of the 3' ORF labeled 'pp' ('proximal protease') encodes the amino terminus of the NS3 protein (residues 1 to 75). 'Bpp' indicates that this region is derived from the HCV Con1 sequence. Both replicons contain the S2204I mutation in NS5A (S→I). 'δ' Indicates the hepatitis delta ribozyme sequence introduced downstream of the 3' terminus of the HCV sequence that produces an exact 3' end.

Figure. 2. Transient HCV RNA replication assay. Shown is the expression of SEAP by En5-3 cells following transfection with the chimeric 1a replicon Bpp-Htat2ANeo/SI and Bpp-Htat2ANeo/KR/SI, which carries an additional K1691R mutation in NS3 that was identified following selection of G418-resistant cells following transfection with Bpp-Htat2ANeo/SI. As controls, SEAP expression is shown following transfection of cells with the highly replication competent 1b replicon, Bpp-Ntat2ANeo/SI, and a related replication defective ΔGDD mutant; also shown in SEAP expression by normal En5-3 cells. Results shown represent the mean values obtained from triplicate cultures transfected with each RNA. SI, S2204 adaptive mutation; KR, K1691R adaptive mutation.

Figure. 3. (A) Schematic depicting the organization of the 5' end of the second ORF in subgenomic chimeric replicons containing most (Bpp-H34A-Ntat2ANeo/SI) or all (Hpp-H34A-Ntat2ANeo/SI) of the H77 genotype 1a NS34A-coding sequence in the background of the genotype 1b Bpp-

Ntat2ANeo/SI. Genotype 1a sequence (H77) is shown as an open box, genotype 1b sequence (Con1 or HCV-N) as a shaded box. 'Bpp' indicates the presence of genotype 1b sequence from the Con1 strain of HCV in the 5' proximal protease coding sequence, whereas 'Hpp' indicates that this sequence is derived from the genotype 1a H77 sequence. Approximate locations are shown for the adaptive mutations Q1067R (Q→R) and G1188R (G→R), identified in G418-resistant cell clones selected following transfection of Hpp-H34A-Ntat2ANeo/SI. (B) SEAP activity present in supernatant culture fluids collected at 24 hr intervals following transfection of En5-3 cells with various chimeric 1a-1b replicons including Bpp-H34A-Ntat2ANeo/SI, Hpp-H34A-Ntat2ANeo/SI, Hpp-H34A-Ntat2ANeo/QR/SI, and Hpp-H34A-Ntat2ANeo/GR/SI. Control cells were transfected with Bpp-Ntat2ANeo/SI and the replication defective ΔGDD mutant. See legend to Fig. 2 for further details. SI, S2204 adaptive mutation; QR, Q1067R adaptive mutation; and GR, G1188R adaptive mutation.

15

Figure 4. Impact of adaptive mutations on replication competence of the subgenomic genotype 1a replicon, Htat2ANeo/SI. (A) Location of various adaptive mutations within the second ORF (derived entirely from the genotype 1a H77 sequence): Q1067R, P1496L (NS3); K1691R (NS4A); and F2080V and S2204I (NS5A). (B) Transient HCV RNA replication assay. SEAP activity in culture supernatants collected at 12-24 hr intervals following electroporation of En5-3 cells with the 1a replicon Htat2ANeo carrying the indicated combinations of the adaptive mutations shown in panel A. Cells were also transfected with genotype 1b Bpp-Ntat2ANeo/SI replicon RNA as a reference. (C) Summary of the replication phenotypes of genotype 1a replicon Htat2ANeo RNAs containing various combinations of adaptive mutations: (-) no detectable replication, (+) modest increase in SEAP expression above background days 3-5, and (+++) >10-fold increase in SEAP expression above background 7 days after transfection in the transient replication assay (see panel B). SI, S2204 adaptive mutation; QR, Q1067R adaptive mutation; PL, P1496L adaptive mutation; KR, K1691R adaptive mutation; and FV, F2080V adaptive mutation.

30

Figure. 5. Adaptive mutations within the polyprotein do not influence the efficiency of polyprotein translation under control of the EMCV IRES. Shown is

an SDS-PAGE gel loaded with products of in vitro translation reactions programmed with RNAs derived from Bpp-Ntat2ANeo (lane 1), Bpp-Htat2ANeo (lanes 2 and 3), Htat2ANeo (lanes 4 to 8), or Bpp-Ntat2ANeo/ Δ GDD (lane 9) RNAs carrying various combinations of adaptive mutations (Q1067R, K1691R, F2080V, or S2204I) as indicated. The schematic at the top of the figure indicates the location of these mutations within the polyprotein. 'pp' indicates the RNA segment encoding the amino terminal 75 residues of NS3, while 'NS' indicates the remainder of the RNA segment encoding the nonstructural proteins. H = genotype 1a H77 sequences, B = genotype 1b Con1 sequences, and N = genotype 1b HCV-N sequences. Location of NS3 and Neo product is indicated at the side of gel.

Figure 6. Impact of additional adaptive mutations on replication competence of the subgenomic genotype 1a replicon, Htat2ANeo/QR/KR/SI (see Fig. 4). (A) Location of various adaptive mutations within the second ORF (derived entirely from the genotype 1a H77 sequence): Q1067R, V1655I (NS3); K1691R (NS4A); and K2040R (KR^{5A}), F2080V and S2204I (NS5A). (B) Transient HCV RNA replication assay. SEAP activity in culture supernatants collected at 12-24 hr intervals following electroporation of En5-3 cells with the 1a replicon Htat2ANeo carrying the indicated combinations of the adaptive mutations shown in panel A. Cells were also transfected with genotype 1b Bpp-Ntat2ANeo/SI replicon RNA as a reference. QR, Q1067R adaptive mutation; VI, V1655I adaptive mutation; KR, K1691R adaptive mutation; KR^{5A}, K2040 adaptive mutation; FV, F2080V adaptive mutation; and SI, S2204I adaptive mutation.

25

Figure 7. Northern analysis of HCV RNA abundance 4 days following transfection of normal Huh7 or En5-3 cells with the indicated dicistronic subgenomic and monocistronic genome length HCV RNAs: (lane 1), normal cells; (lane 2), the subgenomic replicon, Htat2ANeo/SI; (lanes 2-5), Htat2ANeo/SI replicon RNAs carrying the indicated combinations of mutations; (lane 6), nonreplicating Htat2ANeo/QR/VI/KR//KR^{5A}/SI/AAG; (lanes 7) genome-length H77c RNA; (lanes 8-10), genome-length H77c RNA containing the indicated combinations of mutations; (lane 11), genome-length H77 RNA containing the lethal NS5B mutation; (lanes 12 and 13) subcontrol genomic and genome-length

synthetic RNA transcripts. Blots were probed with a genotype 1a probe derived from the NS5B coding sequence for detection of HCV-specific sequence (top panels); blots were also probed for b-actin message to assess RNA loading (lower panels). At the top of the figure is shown the En5-3 cell culture supernatant fluid
5 SEAP activity induced by replicating subgenomic RNAs at the time of cell harvest. SI, S2204 adaptive mutation; QR, Q1067R adaptive mutation; KR, K1691R adaptive mutation; and FV, F2080V adaptive mutation.

Figure 8. Structure of the NS3/4A serine protease/helicase enzyme
10 complex derived from the genotype 1b BK strain of HCV (PDP 1CU1), with the locations of adaptive mutations highlighted. (A) Wire diagram of structure showing the NS3 helicase domain (H) and the protease domain (P). The NS4A cofactor polypeptide (NS4A) is shown in space-filling view, with the NS3 protease active site residues (Active Site) shown in space-filling view. Adaptive
15 mutations identified in this study (Q1067, G1188, V1655, and K1691) cluster near the protease active site or at sites involved in substrate recognition, including the mutations in the NS3 protease domain at Gln-1067, Gly-1188 and near the carboxyl terminus of NS3 in the helicase domain at Val-1655. The NS4A adaptive mutation at Lys-1691 is just beyond the surface of the protease, at the site of exit
20 of the NS4A strand. Adaptive mutations within the NS3 helicase domain that were identified in other studies, S1222, A1226, and P1496 are shown in space-filling view, and are not close to the protease active site. (B) Space-filling view of the structure shown in panel A, in which the adaptive mutations and active site have similar shading. The NS3/4A adaptive mutations identified in this study
25 (Q1067R, G1188R, V1655I, and K1691R) all occur at solvent accessible residues on this side of the molecule. (C) Flip-view of the structure shown in panel B, rotated approximately 180 degrees. The helicase adaptive mutations identified in previous studies are located on the surface of the helicase, distant from the protease active site. Note that in the sequence of the genotype 1b BK strain of
30 HCV, Pro-1496 is Arg (referred to as P1496(R) in the figure, and Lys-1691 is Ser (referred to as K1691(S) in the figure).

Figure 9. Nucleotide sequence of HIVSEAP (SEQ ID NO:7). The HIV long terminal repeat (LTR) is depicted at nucleotides 1-719, and secretory alkaline phosphatase is encoded by the nucleotides 748-2239.

5 Figure 10. 10A, nucleotide sequence of a 3' NTR (SEQ ID NO:8); 10B, nucleotide sequence of a 5' NTR (SEQ ID NO:9).

Figure 11. 11A, nucleotide sequence of a genomic length (full length) hepatitis C virus, genotype 1a (SEQ ID NO:11); 11B, the amino acid sequence of
10 the HCV polyprotein (SEQ ID NO:12) encoded by the coding region present in SEQ ID NO:11.

Figure 12. 12A, nucleotide sequence of Htat2ANeo (SEQ ID NO:13), where nucleotide 1-341 are the 5'NTR, nucleotides 342-1454 are the tat2ANeo
15 (termination codon at 1455-1457), nucleotides 1458-2076 are the EMCV IRES, nucleotides 2080-8034 encode the HCV polyprotein (initiation codon at nucleotides 2077-2079 and termination codon at nucleotides 8035-8037), nucleotides 8038-8259 are the 3'NTR, and nucleotides 8260-8345 are the HDV delta ribozyme (plasmid vector sequences are shown at nucleotides 8346-11240);
20 12B, the amino acid sequence of the HCV polyprotein (SEQ ID NO:14) encoded by the coding region present in SEQ ID NO:13.

Figure 13. Nucleotide (SEQ ID NO:1) of Hepatitis C virus strain H77 and amino acid sequence (SEQ ID NO:2) encoded by nucleotides 342 - 9377.

25

Figure 14. Nucleotide (SEQ ID NO:3) of Hepatitis C virus strain H and amino acid sequence (SEQ ID NO:4) encoded by nucleotides 342 - 9377.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS OF THE INVENTION

30

The present invention provides replication competent polynucleotides. The polynucleotides include a 5' non-translated region (NTR), a 3' NTR, and a coding sequence present between the 5' NTR and 3' NTR. The replication

competent polynucleotides of the present invention are based on hepatitis C virus (HCV), a positive-strand virus. While the ability of a polynucleotide to replicate typically requires the presence of the positive-strand RNA polynucleotide in a cell, it is understood that the term "replication competent polynucleotide" also includes the complement thereof (i.e., the negative-sense RNA), and the corresponding DNA sequences of the positive-sense and the negative-sense RNA sequences. Optionally, a replication competent polynucleotide may be isolated. "Isolated" means a biological material, for instance a polynucleotide, polypeptide, or virus particle, that has been removed from its natural environment. For instance, a virus that has been removed from an animal or from cultured cells in which the virus was propagated is an isolated virus. An isolated polypeptide or polynucleotide means a polypeptide or polynucleotide that has been either removed from its natural environment, produced using recombinant techniques, or chemically or enzymatically synthesized. A "purified" biological material is one that is at least 60% free, preferably 75% free, and most preferably 90% free from other components with which it is naturally associated.

The coding sequence encodes a hepatitis C virus polyprotein. In some aspects of the invention, the HCV polyprotein can yield the following polypeptides; core (also referred to as C or nucleocapsid), E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B. Optionally, a full length HCV polyprotein also yields protein F (see Xu et al., *EMBO J.*, 20, 3840-3848 (2001)). In some aspects of the present invention, an HCV polyprotein is shortened and yields a subset of polypeptides, and typically does not include polypeptides encoded by the amino terminal end of the full length HCV polyprotein. Thus, a hepatitis C virus polyprotein may encode the polypeptides E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B; E2, P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B; P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B; NS2, NS3, NS4A, NS4B, NS5A, and NS5B; or NS3, NS4A, NS4B, NS5A, and NS5B. The hepatitis C virus encoding such a shortened HCV polyprotein may be referred to as a subgenomic hepatitis C virus, and the shortened HCV polyprotein may be referred to as a subgenomic HCV polyprotein. In other aspects of the invention, a replication competent polynucleotide encodes an HCV polyprotein that does not include polypeptides present in an internal portion of a hepatitis C virus polyprotein. Thus, a

subgenomic hepatitis C virus polyprotein may encode, for instance, the polypeptides NS3, NS4A, NS4B, and NS5B.

In those aspects of the invention where the replication competent polynucleotide includes a coding region that encodes less than a full length HCV polyprotein, the 5' end of the coding region encoding the HCV polyprotein may further include about 33 to about 51 nucleotides, or about 36 to about 48 nucleotides, that encode the first about 11 to about 17, or about 12 to about 16, amino acids of the core polypeptide. The result is a fusion polypeptide made up of amino terminal amino acids of the core polypeptide and the first polypeptide encoded by the first cleavage product of the polyprotein, e.g., E1, or E2, or P7, or NS2, etc.

A polyprotein that can yield the core, E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B polypeptides (a full length polyprotein) is typically between about 3000 and 3033 amino acids in length, preferably about 3011 amino acids in length. The relationship between such a polyprotein and the corresponding residues of the individual polypeptides resulting after post-translational processing is shown in Table 1. This numbering system is used herein when referring to a full length polyprotein, and when referring to a polyprotein that contains a portion of the full length polyprotein. For instance, in those aspects of the invention where the replication competent polynucleotide includes a coding sequence encoding an HCV polyprotein that yields the cleavage products NS3, NS4A, NS4B, NS5A, and NS5B and there is no fusion polypeptide made up of amino terminal amino acids of the core polypeptide and the cleavage product NS3, the first amino acid of the NS3 polypeptide is considered to be about residue number 1027. A person of ordinary skill in the art recognizes that this numbering system can vary between members of different genotypes, and between members of the same genotype, thus the numbers shown in Table 1 are approximate, and can vary by 1, 2, 3, 4, or about 5.

Table 1. Correspondence between amino acids of polyprotein and individual polypeptides after processing.

Amino acids of HCV polyprotein ^a	Corresponding polypeptide after processing
1-191	Core
192-383	E1
384-746	E2
747-809	P7
810-1026	NS2
1027-1657	NS3
1658-1711	NS4A
1712-1972	NS4B
1973-2420	NS5A
2421-3011	NS5B

^a Refers to the approximate amino acid number prior to cleavage of the polyprotein where the first amino acid is the first amino acid of the polyprotein expressed by the HCV at Genbank Accession number AF011751 and Genbank Accession number M67463.

A replication competent polynucleotide of the present invention includes at least one adaptive mutation. As used herein, an adaptive mutation is a change in the amino acid sequence of the polyprotein that increases the ability of a replication competent polynucleotide to replicate compared to a replication competent polynucleotide that does not have the adaptive mutation. One adaptive mutation that a replication competent polynucleotide of the present invention typically includes is an isoleucine at about amino acid 2204, which is about amino acid 232 of NS5A. Most clinical HCV isolates and molecularly cloned laboratory HCV strains include a serine at this position, and this mutation has been referred to in the art as S2204I. In most replication competent polynucleotides, the location of this adaptive mutation can also be determined by locating the amino acid sequence SSSA beginning at about amino acid 2200 in the HCV polyprotein, where the amino acid immediately following the SSSA sequence is isoleucine.

A replication competent polynucleotide of the present invention may also include one or more of the adaptive mutations described herein, or a combination thereof. The first such adaptive mutation is an arginine at about amino acid 1067, which is about amino acid 41 of NS3. Most clinical HCV isolates and molecularly cloned laboratory HCV strains include a glutamine at this position, thus this mutation can be referred to as Q1067R. In most replication competent polynucleotides, the location of this adaptive mutation can also be determined by locating the amino acid sequence STAT beginning at about amino acid 1063 in

the HCV polyprotein, where the amino acid immediately following the STAT sequence is arginine. The second adaptive mutation is an arginine at about amino acid 1691, which is about amino acid 34 of NS4A. Most clinical HCV isolates and molecularly cloned laboratory HCV strains include a lysine at this position, thus this mutation can be referred to as K1691R. In most replication competent polynucleotides, the location of this adaptive mutation can also be determined by locating the amino acid sequence VLSG beginning at about amino acid 1687 in the HCV polyprotein, where the amino acid immediately following the VLSG sequence is arginine. The third adaptive mutation is a valine at about amino acid 2080, which is about amino acid 108 of NS5A. Most clinical HCV isolates and molecularly cloned laboratory HCV strains include a phenylalanine at this position, thus this mutation can be referred to as F2080V. In most replication competent polynucleotides, the location of this adaptive mutation can also be determined by locating the amino acid sequence ALWR beginning at about amino acid 2081 in the HCV polyprotein, where the amino acid immediately before the ALWR sequence is valine. A fourth adaptive mutation is an isoleucine at about amino acid 1655, which is about amino acid 629 of NS3. Most clinical HCV isolates and molecularly cloned laboratory HCV strains include a valine at this position, thus this mutation can be referred to as V1655I. In most replication competent polynucleotides, the location of this adaptive mutation can also be determined by locating the amino acid sequence ADLE beginning at about amino acid 2051 in the HCV polyprotein, where the amino acid immediately after the ADLE sequence is isoleucine. A fifth adaptive mutation is an arginine at about amino acid 2040, which is about amino acid 68 of NS5A. Most clinical HCV isolates and molecularly cloned laboratory HCV strains include a lysine at this position, thus this mutation can be referred to as K2040R. In most replication competent polynucleotides, the location of this adaptive mutation can also be determined by locating the amino acid sequence GHVXN beginning at about amino acid 2037 in the HCV polyprotein, where the X in the amino acid is arginine. A sixth adaptive mutation is an arginine at about amino acid 1188, which is about amino acid 162 of NS3. Most clinical HCV isolates and molecularly cloned laboratory HCV strains include a glycine at this position, thus this mutation can be referred to as G1188R. In most replication competent polynucleotides, the location of this adaptive mutation can also be determined by

locating the amino acid sequence VCTR beginning at about amino acid 1184 in the HCV polyprotein. In some aspects, the replication competent polynucleotide of the present invention includes the Q1067R and K1691R adaptive mutations, as well as the S2204I adaptive mutation. These adaptive mutations are summarized in Table 2. A person of ordinary skill in the art recognizes that the precise location of these cell culture adaptive mutations can vary between members of different genotypes, and between members of the same genotype, thus the numbers shown in Table 2 are approximate, and can vary by 1, 2, 3, 4, or about 5.

Table 2. Adaptive Mutations

Symbol ¹	Protein / Residue ²	Mutation ³
QR	NS3/41	Q1067R
GR	NS3/162	G1188R
VI	NS3/629	V1655I
KR	NS4A/34	K1691R
KR ^{5A}	NS5A/68	K2040R
FV	NS5A/108	F2080V
SI	NS5A/232	S2204I

¹Symbol used to designate presence in RNA transcripts.

²Residue refers to position in protein after post-translational cleavage of the H77c polyprotein (GenBank accession AF011751).

³Number refers to position of mutation in H77c polyprotein before post-translational cleavage (GenBank accession AF011751).

There are many other adaptive mutations known to the art, and the replication competent polynucleotides of the present invention may include one or more of those adaptive mutations. Examples of known adaptive mutations can be found in, for instance, Bartenschlager (U.S. Patent 6,630,343), Blight et al. (Science, 290, 1972-1975 (2000)), Lohmann et al., (Abstract P038, 7th International Meeting on Hepatitis C virus and Related viruses (Molecular Virology and Pathogenesis), December 3-7 (2000)), Guo et al. (Abstract P045, 7th International Meeting on Hepatitis C virus and Related viruses (Molecular

Virology and Pathogenesis), December 3-7 (2000)), Blight et al., (J. Virol. 77, 3181-3190 (2003)), Gu et al., (J. Virol. 77, 5352-5359 (2003)), and Grobler et al., (J. Biol. Chem., 278,16741-16746 (Feb, 2003).

It is expected that polynucleotides encoding an HCV polyprotein can be
5 obtained from different sources, including molecularly cloned laboratory strains, for instance cDNA clones of HCV, and clinical isolates. Examples of molecularly cloned laboratory strains include the HCV that is encoded by pCV-H77C (Yanagi et al., *Proc. Natl. Acad. Sci. USA*, 94, 8738-8743 (1997), Genbank accession number AF011751, SEQ ID NO:1), and pHCV-H (Inchauspe et al., *Proc. Natl.*
10 *Acad. Sci. USA*, 88, 10292-10296 (1991), Genbank accession number M67463, SEQ ID NO:3). Clinical isolates can be from a source of infectious HCV, including tissue samples, for instance from blood, plasma, serum, liver biopsy, or leukocytes, from an infected animal, including a human or a primate. It is also expected that the polynucleotide encoding the HCV polyprotein present in a
15 replication competent polynucleotide can be prepared by recombinant, enzymatic, or chemical techniques. The nucleotide sequence of molecularly cloned laboratory strains and clinical isolates can be modified to encode an HCV polyprotein that includes the S2204I adaptive mutation and one or more of the adaptive mutations described herein. Such methods are routine and known to the
20 art and include, for instance, PCR mutagenesis.

The present invention further includes replication competent polynucleotides encoding an HCV polyprotein having similarity with the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4 (in the case of a full length polyprotein), or a portion thereof (in the case of an HCV polyprotein encoding, for
25 instance, NS3, NS4A, NS4B, NS5A, and NS5B, and not encoding core, E1, E2, P7, and NS2). The similarity is referred to as structural similarity and is generally determined by aligning the residues of the two amino acid sequences (i.e., a candidate amino acid sequence and the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, or a portion thereof) to optimize the number of identical amino
30 acids along the lengths of their sequences; gaps in either or both sequences are permitted in making the alignment in order to optimize the number of identical amino acids, although the amino acids in each sequence must nonetheless remain in their proper order. A candidate amino acid sequence is the amino acid sequence being compared to an amino acid sequence present in SEQ ID NO:2,

SEQ ID NO:4, or a portion thereof. A candidate amino acid sequence can be isolated from a cell infected with a hepatitis C virus, or can be produced using recombinant techniques, or chemically or enzymatically synthesized. Preferably, two amino acid sequences are compared using the Blastp program of the BLAST
5 2 search algorithm, as described by Tatusova, et al. (*FEMS Microbiol Lett* 1999, 174:247-250), and available at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. Preferably, the default values for all BLAST 2 search parameters are used, including matrix = BLOSUM62; open gap penalty = 11, extension gap penalty = 1, gap x_dropoff = 50, expect = 10, wordsize = 3, and optionally, filter on. In the
10 comparison of two amino acid sequences using the BLAST search algorithm, structural similarity is referred to as "identities." An HCV polyprotein may include an amino acid sequence having a structural similarity with SEQ ID NO:2, SEQ ID NO:4, or a portion thereof, of at least about 90 %, for example 91%, 92%, 93% identity, and so on to 100 % identity. A replication competent
15 polynucleotide having a 5' NTR of SEQ ID NO:9, a 3' NTR of SEQ ID NO:8, and HCV polyprotein with structural similarity with SEQ ID NO:2, SEQ ID NO:4, or a portion thereof, is replication competent in a cell derived from a human hepatoma such as Huh-7 and Huh-7.5. An HCV polyprotein having structural similarity with the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, or a
20 portion thereof, includes the S2204I adaptive mutation and one or more of the adaptive mutations described herein. Such an HCV polyprotein may optionally include other adaptive mutations.

In some aspects, the coding sequence of a replication competent polynucleotide of the present invention that encodes a hepatitis C virus
25 polyprotein is not a specific genotype. For instance, a polynucleotide encoding an HCV polyprotein present in a replication competent polynucleotide of the present invention can be genotype 1a, 1b, 1c, 2a, 2b, 2c, 3a, 3b, 4, 5a, or 6a (as defined by Simmonds, *Hepatology*, 21, 570-583 (1995)). In other aspects, the HCV polyprotein is genotype 1a. Methods for determining the genotype of a hepatitis C
30 virus are routine and known to the art and include, for instance, serotyping the virus particle using antibody, and/or evaluation of the nucleotide sequence by, for instance, polymerase chain reaction assays (see Simmonds, *J. Hepatol.*, 31(Suppl. 1), 54-60 (1999)).

The present invention includes polynucleotides encoding an amino acid sequence having similarity to an HCV polyprotein. The similarity is referred to as structural similarity and is determined by aligning the residues of two polynucleotides (e.g., the nucleotide sequence of the candidate coding region and nucleotides 342 - 9377 of SEQ ID NO:1 or nucleotides 342 - 9377 of SEQ ID NO:3) to optimize the number of identical nucleotides along the lengths of their sequences; gaps in either or both sequences are permitted in making the alignment in order to optimize the number of shared nucleotides, although the nucleotides in each sequence must nonetheless remain in their proper order. A candidate coding region is the coding region being compared to a coding region present in SEQ ID NO:1 (e.g., nucleotides 342 - 9377 of SEQ ID NO:1). A candidate nucleotide sequence can be isolated from a cell, or can be produced using recombinant techniques, or chemically or enzymatically synthesized. Preferably, two nucleotide sequences are compared using the Blastn program of the BLAST 2 search algorithm, as described by Tatusova, et al. (*FEMS Microbiol Lett* 1999, 174:247-250), and available at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. Preferably, the default values for all BLAST 2 search parameters are used, including reward for match = 1, penalty for mismatch = -2, open gap penalty = 5, extension gap penalty = 2, gap x_dropoff = 50, expect = 10, wordsize = 11, and optionally, filter on. In the comparison of two nucleotide sequences using the BLAST search algorithm, structural similarity is referred to as "identities."

The present invention also includes polynucleotides encoding the HCV polyproteins described herein, including, for instance, the polyproteins having the amino acid sequence shown in SEQ ID NO:2 and SEQ ID NO:4. An example of the class of nucleotide sequences encoding each of these polyproteins are nucleotides 342 - 9377 of SEQ ID NO:1 and nucleotides 342 - 9377 of SEQ ID NO:3, respectively. These classes of nucleotide sequences are large but finite, and the nucleotide sequence of each member of the class can be readily determined by one skilled in the art by reference to the standard genetic code.

A replication competent polynucleotide of the present invention includes a 5' non-translated region (NTR) (see Smith et al., *J. Gen. Virol.*, 76, 1749-1761 (1995)). A 5' NTR is typically about 341 nucleotides in length. A replication competent polynucleotide of the present invention also includes a 3' NTR. A 3' NTR typically includes, from 5' to 3', nucleotides of variable length and sequence

(referred to as the variable region), a poly-pyrimidine tract (the poly U-UC region), and a highly conserved sequence of about 100 nucleotides (the conserved region) (see, for instance, Lemon et al., U.S. Published Application US 2003 0125541, and Yi and Lemon, *J. Virol.*, 77, 3557-3568 (2003)). The variable
5 region begins at about the first nucleotide following the stop codon of the HCV polyprotein, and generally ends immediately before the nucleotides of the poly U-UC region. The poly U-UC region is a stretch of predominantly U residues, CU residues, or C(U)_n-repeats. When the nucleotide sequence of a variable region is compared between members of the same genotype, there is typically a great deal
10 of similarity; however, there is typically very little similarity in the nucleotide sequence of the variable regions between members of different genotypes (see, for instance, Yamada et al., *Virology*, 223, 255-261 (1996)).

It is expected that a 5' NTR and a 3' NTR can be obtained from different sources, including molecularly cloned laboratory strains, for instance cDNA
15 clones of HCV, and clinical isolates. Examples of molecularly cloned laboratory strains include the HCV that is encoded by pCV-H77C (Yanagi et al., *Proc. Natl. Acad. Sci. USA*, 94, 8738-8743 (1997), Genbank accession number AF011751, SEQ ID NO:1, where nucleotides 1-341 are the 5' NTR and nucleotides 9378-9599 are the 3' NTR), and pHCV-H (Inchauspe et al., *Proc. Natl. Acad. Sci. USA*,
20 88, 10292-10296 (1991), Genbank accession number M67463, SEQ ID NO:3, where nucleotides 1-341 are the 5' NTR and nucleotides 9378-9416 are the 3' NTR). Clinical isolates can be from a source of infectious HCV, including tissue samples, for instance from blood, plasma, serum, liver biopsy, or leukocytes, from an infected animal, including a human or a primate. It is also expected that the
25 polynucleotide encoding the HCV polyprotein present in a replication competent polynucleotide can be prepared by recombinant, enzymatic, or chemical techniques.

In some aspects, a 5' NTR and a 3' NTR of a replication competent polynucleotide of the present invention is not a specific genotype. For instance, a
30 5' NTR and a 3' NTR present in a replication competent polynucleotide of the present invention can be genotype 1a, 1b, 1c, 2a, 2b, 2c, 3a, 3b, 4, 5a, or 6a (as defined by Simmons, *Hepatology*, 21, 570-583 (1995)). In other aspects, the HCV polyprotein is genotype 1a. Methods for determining the genotype of a 5' NTR and a 3' NTR are routine and known to the art and include evaluation of the

nucleotide sequence for specific nucleotides that are characteristic of a specific genotype.

In some aspects of the invention a replication competent polynucleotide includes a second coding region. The second coding sequence may be present in the 3' NTR, for instance, in the variable region of the 3' NTR. In some aspects of the invention, the second coding region is present in the variable region such that the variable region is not removed. Alternatively, the second coding region replaces the variable region in whole or in part. In some aspects of the invention, for instance, when the HCV has the genotype 1a, the second coding region is inserted in the variable region between nucleotides 5 and 6 of the sequence 5' CUCUUAAGC 3', where the sequence shown corresponds to the positive-strand.

In some aspects of the invention, the second coding region is present in a replication competent polynucleotide downstream of the 5' NTR, and upstream of the first coding region, i.e., the coding region encoding a HCV polyprotein. For instance, the first nucleotide of the second coding region may be immediately downstream and adjacent to the last nucleotide of the 5' NTR. Alternatively, the first nucleotide of the second coding region may be further downstream of the last nucleotide of the 5' NTR, for instance, about 2 to about 51 nucleotides, about 33 to about 51 nucleotides, or about 36 to about 48 nucleotides downstream of the last nucleotide of the 5' NTR. Typically, when the first nucleotide of the second coding region is not immediately downstream of the last nucleotide of the 5' NTR, the nucleotides in between the 5' NTR and the second coding region encode the amino terminal amino acids of the HCV core polypeptide. For instance, the 5' end of the second coding region may further include about 33 to about 51 nucleotides, or about 36 to about 48 nucleotides, that encode the first about 11 to about 17, or about 12 to about 16, amino acids of the core polypeptide. The result is a fusion polypeptide made up of amino terminal amino acids of the core polypeptide and the polypeptide encoded by the second coding region (see, for instance, Yi et al., Virol., 304, 197-210 (2002), and U.S. Published Application US 2003 0125541). Without intending to be limiting, it is believed the presence of the nucleotides from the core coding sequence act to enhance translation the polypeptide encoded by the second coding region.

In those aspects of the invention where the second coding region present in a replication competent polynucleotide is present downstream of the 5' NTR and

upstream of the coding region encoding the HCV polyprotein, the replication competent polynucleotide typically includes a regulatory region operably linked to the downstream coding region, e.g., the coding region encoding the HCV polyprotein. Preferably, the regulatory region provides for the translation of the downstream coding region. The size of the regulatory region may be from about 400 nucleotides to about 800 nucleotide, more preferably, about 600 nucleotides to about 700 nucleotides. Typically, the regulatory region is an IRES. Examples of IRES elements are described herein.

The second coding region can encode a polypeptide including, for instance, a marker, including a detectable marker and/or a selectable marker. Examples of detectable markers include molecules having a detectable enzymatic activity, for instance, secretory alkaline phosphatase, molecules having a detectable fluorescence, for instance, green or red or blue fluorescent protein, and molecules that can be detected by antibody. Examples of selectable markers include molecules that confer resistance to antibiotics able to inhibit the replication of eukaryotic cells, including the antibiotics kanamycin, ampicillin, chloramphenicol, tetracycline, blasticidin, neomycin, and formulations of phleomycin D1 including, for example, the formulation available under the trade-name ZEOCIN (Invitrogen, Carlsbad, California). Coding sequences encoding such markers are known to the art. Other examples of polypeptides that can be encoded by the second coding region include a transactivator, and/or a fusion polypeptide. Preferably, when the polypeptide is a fusion polypeptide, the second coding region includes nucleotides encoding a marker, more preferably, nucleotides encoding a fusion between a transactivator and a marker. Transactivators are described herein below. Optionally, the coding region can encode an immunogenic polypeptide. A replication competent polynucleotide containing a second coding region is typically dicistronic, i.e., the coding region encoding the HCV polyprotein and the second coding region are separate.

An "immunogenic polypeptide" refers to a polypeptide which elicits an immunological response in an animal. An immunological response to a polypeptide is the development in a subject of a cellular and/or antibody-mediated immune response to the polypeptide. Usually, an immunological response includes but is not limited to one or more of the following effects: the production

of antibodies, B cells, helper T cells, suppressor T cells, and/or cytotoxic T cells, directed specifically to an epitope or epitopes of the polypeptide fragment.

A transactivator is a polypeptide that affects in *trans* the expression of a coding region, preferably a coding region integrated in the genomic DNA of a cell. Such coding regions are referred to herein as "transactivated coding regions." The cells containing transactivated coding regions are described in detail herein below. Transactivators useful in the present invention include those that can interact with a regulatory region, preferably an operator sequence, that is operably linked to a transactivated coding region. As used herein, the term "transactivator" includes polypeptides that interact with an operator sequence and either prevent transcription from initiating at, activate transcription initiation from, or stabilize a transcript from, a transactivated coding region operably linked to the operator sequence. Examples of useful transactivators include the HIV tat polypeptide (see, for example, the polypeptides

MEPVDPRLEPWKHPGSQPKTACTNCYCKKCCFHCQVCFITKALGISYGRK
KRRQRRRAHQNSQTHQASLSKQPTSQPRGDPTGPKE (SEQ ID NO:5) which is encoded by nucleotides 5377 to 5591 and 7925 to 7970 of Genbank accession number AF033819), and

MEPVDPRLEPWKHPGSQPKTACTNCYCKKCCFHCQVCFITKALGISYGRK
KRRQRRRPPQGSQTHQVSLSKQPTSQSRGDPTGPKE (SEQ ID NO:10). The HIV tat polypeptide interacts with the HIV long terminal repeat (LTR). Other useful transactivators include human T cell leukemia virus tax polypeptide (which binds to the operator sequence tax response element, Fujisawa et al., *J. Virol.*, 65, 4525-4528 (1991)), and transactivating polypeptides encoded by spumaviruses in the region between env and the LTR, such as the bel-1 polypeptide in the case of human foamy virus (which binds to the U3 domain of these viruses, Rethwilm et al., *Proc. Natl. Acad. Sci. USA*, 88, 941-945 (1991)). Alternatively, a post-transcriptional transactivator, such as HIV rev, can be used. HIV rev binds to a 234 nucleotide RNA sequence in the env gene (the rev-response element, or RRE) of HIV (Hadzopolou-Cladaras et al., *J. Virol.*, 63, 1265-1274 (1989)).

Other transactivators that can be used are those having similarity with the amino acid sequence of SEQ ID NO:5 or SEQ ID NO:10. The similarity is generally determined as described herein above. A candidate amino acid sequence that is being compared to an amino acid sequence present in SEQ ID

NO:5 or SEQ ID NO:10 can be isolated from a virus, or can be produced using recombinant techniques, or chemically or enzymatically synthesized. Preferably, two amino acid sequences are compared using the Blastp program of the BLAST 2 search algorithm, as described herein above. Preferably, a transactivator

5 includes an amino acid sequence having a structural similarity with SEQ ID NO:5 or SEQ ID NO:10, of at least about 90 %, at least about 94 %, at least about 96 %, at least about 97 %, at least about 98 %, or at least about 99 % identity. Typically, an amino acid sequence having a structural similarity with SEQ ID NO: 5 or SEQ ID NO:10 has tat activity. Whether such a polypeptide has activity can be

10 evaluated by determining if the amino acid sequence can interact with an HIV LTR, preferably alter transcription from a coding sequence operably linked to an HIV LTR. Useful HIV LTRs are described herein.

Active analogs or active fragments of a transactivator can be used in the invention. An active analog or active fragment of a transactivator is one that is

15 able to interact with an operator sequence and either prevent transcription from initiating at, activate transcription initiation from, or stabilize a transcript from, a transactivated coding region operably linked to the operator sequence.

Active analogs of a transactivator include polypeptides having conservative amino acid substitutions that do not eliminate the ability to interact

20 with an operator and alter transcription. Substitutes for an amino acid may be selected from other members of the class to which the amino acid belongs. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and tyrosine. Polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, aspartate, and glutamate.

25 The positively charged (basic) amino acids include arginine, lysine, and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Examples of preferred conservative substitutions include Lys for Arg and *vice versa* to maintain a positive charge; Glu for Asp and *vice versa* to maintain a negative charge; Ser for Thr so that a free -OH is maintained; and Gln for Asn to

30 maintain a free NH₂.

Active fragments of a transactivator include a portion of the transactivator containing deletions or additions of about 1, about 2, about 3, about 4, or at least about 5 contiguous or noncontiguous amino acids such that the resulting transactivator will alter expression of an operably linked transactivated coding

region. A preferred example of an active fragment of the HIV tat polypeptide includes amino acids 1-48 of SEQ ID NO: 5, or amino acids 1-48 of SEQ ID NO:10.

In those aspects of the invention where the second coding region encodes a fusion polypeptide, the fusion polypeptide can further include amino acids corresponding to a *cis*-active proteinase. When the fusion polypeptide is a fusion between a transactivator and a marker, preferably the fusion polypeptide also includes amino acids corresponding to a *cis*-active proteinase. Preferably the amino acids corresponding to a *cis*-active proteinase are present between the amino acids corresponding to the transactivator and the marker. A *cis*-active proteinase in this position allows the amino acids corresponding to the transactivator and the marker to be physically separate from each other in the cell within which the replication competent polynucleotide is present. Examples of *cis*-active proteinases that are useful in the present invention include the *cis*-active 2A proteinase of foot-and-mouth disease (FMDV) virus (see, for example, US Patent 5,846,767 (Halpin et al.) and US Patent 5,912,167 (Palmenberg et al.)), ubiquitin (see, for example, Tautz et al., *Virology*, 197, 74-85 (1993)), and the NS3 recognition site GADTEDVVCCSMSY (SEQ ID NO:6) (see, for example, Lai et al., *J. Virol.*, 74, 6339-6347 (2000)).

Active analogs and active fragments of *cis*-active proteinases can also be used. Active analogs of a *cis*-acting proteinase include polypeptides having conservative amino acid substitutions that do not eliminate the ability of the proteinase to catalyze cleavage. Active fragments of a *cis*-active proteinase include a portion of the *cis*-active proteinase containing deletions or additions of one or more contiguous or noncontiguous amino acids such that the resulting *cis*-active proteinase will catalyze the cleavage of the proteinase.

In some aspects of the invention, the second coding region may further include an operably linked regulatory region. Preferably, a regulatory region located 5' of the operably linked coding region provides for the translation of the coding region.

A preferred regulatory region located 5' of an operably linked second coding region is an internal ribosome entry site (IRES). An IRES allows a ribosome access to mRNA without a requirement for cap recognition and subsequent scanning to the initiator AUG (Pelletier, et al., *Nature*, 334, 320-325

(1988)). An IRES is located upstream of the translation initiation codon, e.g., ATG or AUG, of the coding sequence to which the IRES is operably linked. The distance between the IRES and the initiation codon is dependent on the type of IRES used, and is known to the art. For instance, poliovirus IRES initiates a
5 ribosome translocation/scanning process to a downstream AUG codon. For other IRES elements, the initiator codon is generally located at the 3' end of the IRES sequence. Examples of an IRES that can be used in the invention include a viral IRES, preferably a picornaviral IRES or a flaviviral IRES. Examples of poliovirus IRES elements include, for instance, poliovirus IRES,
10 encephalomyocarditis virus IRES, or hepatitis A virus IRES. Examples of preferred flaviviral IRES elements include hepatitis C virus IRES, GB virus B IRES, or a pestivirus IRES, including but not limited to bovine viral diarrhea virus IRES or classical swine fever virus IRES. Other IRES elements with similar secondary and tertiary structure and translation initiation activity can either be
15 generated by mutation of these viral sequences, by cloning of analogous sequences from other viruses (including picornaviruses), or prepared by enzymatic synthesis techniques.

The size of the second coding region is not critical to the invention. It is expected there is no lower limit on the size of the second coding region, and that
20 there is an upper limit on the size of the second coding region. This upper limit can be easily determined by a person skilled in the art, as second coding region that are greater than this upper limit adversely affect replication of a replication competent polynucleotide. The second coding region is typically at least about 10 nucleotides, at least about 20 nucleotides, at least about 30 nucleotides, or at least
25 about 40 nucleotides.

A replication competent polynucleotide may also include a nucleotide sequence having cis-acting ribozyme activity. Such a ribozyme is typically present at the 3' end of the 3' NTR of a replication competent polynucleotide, and generates a precise 3' terminal end of the replication competent polynucleotide
30 when it is an RNA molecule by cleaving the junction between the replication competent polynucleotide and the ribozyme. This can be advantageous when the replication competent polynucleotide is to be used for a transient transfection. Since the ribozyme catalyzes its own removal from the RNA molecule, this type

of ribozyme is present only when a replication competent polynucleotide is a DNA molecule.

The replication competent polynucleotide of the invention can be present in a vector. When a replication competent polynucleotide is present in a vector the polynucleotide is DNA, including the 5' non-translated RNA and the 3' non-translated RNA, and, if present, the second coding sequence. Methods for cloning and/or inserting hepatitis C virus sequences into a vector are known to the art (see, e.g., Yanagi et al., *Proc. Natl. Acad. Sci., USA*, 94, 8738-8743 (1997); and Rice et al., (U.S. Patent 6,127,116)). Such constructs are often referred to as molecularly cloned laboratory strains, and an HCV that is inserted into a vector is often referred to as a cDNA clone of the HCV. If the RNA encoded by the HCV is able to replicate *in vivo*, the HCV present in the vector is referred to as an infectious cDNA clone. A vector is a replicating polynucleotide, such as a plasmid, phage, cosmid, or artificial chromosome to which another polynucleotide may be attached so as to bring about the replication of the attached polynucleotide. A vector can provide for further cloning (amplification of the polynucleotide), i.e., a cloning vector, or for expression of the polypeptide encoded by the coding region, i.e., an expression vector. The term vector includes, but is not limited to, plasmid vectors, viral vectors, cosmid vectors, or artificial chromosome vectors. Preferably the vector is a plasmid. Preferably the vector is able to replicate in a prokaryotic host cell, for instance *Escherichia coli*. Preferably, the vector can integrate in the genomic DNA of a eukaryotic cell.

An expression vector optionally includes regulatory sequences operably linked to the replication competent polynucleotide such that it is transcribed to produce RNA molecules. These RNA molecules can be used, for instance, for introducing a replication competent polynucleotide into a cell that is in an animal or growing in culture. The terms "introduce" and "introducing" refer to providing a replication competent polynucleotide to a cell under conditions that the polynucleotide is taken up by the cell in such a way that it can then replicate. The replication competent polynucleotide can be present in a virus particle, or can be a nucleic acid molecule, for instance, RNA. The invention is not limited by the use of any particular promoter, and a wide variety are known. Promoters act as regulatory signals that bind RNA polymerase in a cell to initiate transcription of a downstream (3' direction) HCV. The promoter used in the invention can be a

constitutive or an inducible promoter. A preferred promoter for the production of replication competent polynucleotide as an RNA molecule is a T7 promoter.

The present invention includes methods for identifying a replication competent polynucleotide, including detecting and/or selecting for cells
5 containing a replication competent polynucleotide. Typically, the cells used in this aspect of the invention are primate or human cells growing in culture. Useful cultured cells will support the replication of the polynucleotides of the present invention, and include primary human or chimpanzee hepatocytes, peripheral mononuclear cells, cultured human lymphoid cell lines (for instance lines
10 expressing B-cell and T-cell markers such as Bjab and Molt-4 cells), and continuous cell lines derived from such cells, including HPBMa10-2 and Daudi (Shimizu et al., *J. Gen. Virol.*, 79, 1383-1386 (1998), and MT-2 (Kato et al., *Biochem. Biophys. Res. Commun.*, 206, 863-869 (1995)). Other useful cells include those derived from a human hepatoma cells, for instance, Huh-7 (see, for
15 instance, Lohmann et al. (*Science*, 285, 110-113 (1999)), Huh-7.5 (see, for instance, Blight et al., *J. Virol.*, 76, 13001-13014 (2002), and Blight et al., *J. Virol.*, 77, 3183-3190 (2003)), HepG2 and IMY-N9 (Date et al., *J. Biol. Chem.*, 279, 22371-22376 (2004)), and PH5CH8 (Ikeda et al., *Virus Res.*, 56, 157-167 (1998)). In general, useful cells include those that support replication of HCV RNA,
20 including, for instance, replication of the HCV encoded by pCV-H77C, replication of the HCV encoded by pHCV-N as modified by Beard et al. (*Hepatol.*, 30, 316-324 (1999)), or replication of such an HCV modified to contain one or more adaptive mutations.

In some aspects of the invention, the cultured cell includes a
25 polynucleotide that includes a coding region, the expression of which is controlled by a transactivator. Such a coding region is referred to herein as a transactivated coding region. A transactivated coding region encodes a marker, such as a detectable marker, for example, secretory alkaline phosphatase (SEAP), an example of which is encoded by nucleotides 748-2239 of SEQ ID NO:7 (see Fig.
30 9). Typically, a cultured cell that includes a polynucleotide having a transactivated coding region is used in conjunction with a replication competent polynucleotide of the present invention that includes a coding region encoding a transactivator.

The polynucleotide that includes the transactivated coding region can be present integrated into the genomic DNA of the cell, or present as part of a vector that is not integrated. Methods of modifying a cell to contain an integrated DNA are known to the art (see, for instance, Lemon et al., U.S. Published Application
5 US 2003 0125541, and Yi et al., *Viol.*, 302, 197-210 (2002)).

Operably linked to the transactivated coding region is an operator sequence. The interaction of a transactivator with an operator sequence can alter transcription of the operably linked transactivated coding region. In those aspects of the invention where a transactivator increases transcription, there is typically
10 low transcription, or, essentially no transcription, of the transactivated coding region in the absence of a transactivator. An operator sequence can be present upstream (5') or downstream (3') of a transactivated coding region. An operator sequence can be a promoter, or can be a nucleotide sequence that is present in addition to a promoter.

15 In some aspects of the invention, the operator sequence that is operably linked to a transactivated coding sequence is an HIV long terminal repeat (LTR). An example of an HIV LTR is depicted at nucleotides 1-719 of SEQ ID NO:7. Also included in the present invention are operator sequences having similarity to nucleotides 1-719 of SEQ ID NO:7. The similarity between two nucleotides
20 sequences may be determined by aligning the residues of the two polynucleotides (i.e., the nucleotide sequence of the candidate operator sequence and the nucleotide sequence of nucleotides 1-719 of SEQ ID NO:7) to optimize the number of identical nucleotides along the lengths of their sequences; gaps in either or both sequences are permitted in making the alignment in order to optimize the
25 number of shared nucleotides, although the nucleotides in each sequence must nonetheless remain in their proper order. A candidate operator sequence can be isolated from a cell, or can be produced using recombinant techniques, or chemically or enzymatically synthesized. Preferably, two nucleotide sequences are compared using the Blastn program of the BLAST 2 search algorithm, as
30 described by Tatusova, et al. (*FEMS Microbiol Lett* 1999, 174:247-250), and available at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. Preferably, the default values for all BLAST 2 search parameters are used, including reward for match = 1, penalty for mismatch = -2, open gap penalty = 5, extension gap penalty = 2, gap x_dropoff = 50, expect = 10, wordsize = 11, and filter on. In the comparison of

two nucleotide sequences using the BLAST search algorithm, structural similarity is referred to as "identities." Preferably, an operator sequence includes a nucleotide sequence having a structural similarity with the nucleotides 1-719 of SEQ ID NO:7 of at least about 90 %, at least about 95 %, or at least about 99 % identity. Typically, an operator sequence having structural similarity with the nucleotides 1-719 of SEQ ID NO:7 has transcriptional activity. Whether such an operator sequence has transcriptional activity can be determined by evaluating the ability of the operator sequence to alter transcription of an operably linked coding sequence in response to the presence of a polypeptide having tat activity, preferably, a polypeptide including the amino acids of SEQ ID NO:5 or SEQ ID NO:10.

A selecting agent may be used to inhibit the replication of cultured cells that support the replication of polynucleotides of the present invention. Examples of selecting agents include antibiotics, including kanamycin, ampicillin, chloramphenicol, tetracycline, neomycin, and formulations of phleomycin D1. A selecting agent can act to prevent replication of a cell, or kill a cell, while the agent is present and the cell does not express a molecule that provides resistance to the selecting agent. Typically, the molecule providing resistance to a selecting agent is expressed in the cell by a replication competent polynucleotide of the present invention. Alternatively, the molecule providing resistance to a selecting agent is expressed by the cell but the expression of the molecule is controlled by a replication competent polynucleotide of the present invention that is present in the cell. The concentration of the selecting agent is typically chosen such that a cell does not replicate if it does not contain a molecule providing resistance to a selecting agent. The appropriate concentration of a selecting agent varies depending on the particular selecting agent, and can be easily determined by one having ordinary skill in the art using known techniques.

When a polynucleotide is introduced into a cell that is growing in culture, the polynucleotide can be introduced using techniques known to the art. Such techniques include, for instance, liposome and non-liposome mediated transfection. Non-liposome mediated transfection methods include, for instance, electroporation.

In some aspects of the invention, when a replication competent polynucleotide is identified using cultured cells, its ability to replicate may be

verified by introducing the replication competent polynucleotide into a cell present in an animal, preferably a chimpanzee. When the cell is present in the body of an animal, the replication competent polynucleotide can be introduced by, for instance, subcutaneous, intramuscular, intraperitoneal, intravenous, or percutaneous intrahepatic administration, preferably by percutaneous intrahepatic administration. Methods for determining whether a replication competent polynucleotide is able to replicate in a chimpanzee are known to the art (see, for example, Yanagi et al., *Proc. Natl. Acad. Sci. USA*, 94, 8738-8743 (1997)). In general, the demonstration of infectivity is based on the appearance of the virus in the circulation of the chimpanzee over the days and weeks following the intrahepatic injection of the replication competent polynucleotide. The presence of the virus can be confirmed by reverse transcription-polymerase chain reaction (RT-PCR) detection of the viral RNA, by inoculation of a second chimpanzee with transfer of the hepatitis C virus infection as indicated by the appearance of liver disease and seroconversion to hepatitis C virus in ELISA tests, or possibly by the immunologic detection of components of the hepatitis C virus (e.g., the core protein) in the circulation of the inoculated animal. It should be noted that seroconversion by itself is generally not a useful indicator of infection in an animal injected with a viral RNA produced using a molecularly cloned laboratory strain, as this RNA may have immunizing properties and be capable of inducing HCV-specific antibodies to proteins translated from an input RNA that is non-replicating. Similarly, the absence of seroconversion does not exclude the possibility of viral replication and infection of a chimpanzee with HCV.

Whether a polynucleotide is replication competent can be determined using methods known to the art, including methods that use nucleic acid amplification to detect the result of increased levels of replication. For instance, transient transfection of a cell with a replication competent polynucleotide permits measurement of the production of additional polynucleotides. Methods for transient transfection of a cell with a replication competent polynucleotide and for assay of subsequent replication are known to the art. In some aspects of the invention, another method for detecting a replication competent polynucleotide includes measuring the production of viral particles by a cell. The measurement of viral particles can be accomplished by passage of supernatant from media containing a cell culture that may contain a replication competent polynucleotide,

and using the supernatant to infect a second cell. Detection of the polynucleotide or viral particles in the second cell indicates the initial cell contains a replication competent polynucleotide. The production of infectious virus particles by a cell can also be measured using antibody that specifically binds to an HCV viral
5 particle. As used herein, an antibody that can "specifically bind" an HCV viral particle is an antibody that interacts only with the epitope of the antigen (e.g., the viral particle or a polypeptide that makes up the particle) that induced the synthesis of the antibody, or interacts with a structurally related epitope. "Epitope" refers to the site on an antigen to which specific B cells and/or T cells
10 respond so that antibody is produced. An epitope could include about 3 amino acids in a spatial conformation which is unique to the epitope. Generally an epitope includes at least about 5 such amino acids, and more usually, consists of at least about 8-10 such amino acids. Antibodies to HCV viral particles can be produced as described herein.

15 In another aspect, identifying a replication competent polynucleotide includes incubating a cultured cell that includes a polynucleotide of the present invention. In those aspects of the invention where the replication competent polynucleotide includes a second coding region encoding a detectable marker, cells containing the replication competent polynucleotide can be identified by
20 observing individual cells that contain the detectable marker. Alternatively, if the detectable marker is secreted by the cell, the presence of the marker in the medium in which the cell is incubated can be detected. Methods for observing the presence or absence of a detectable marker in a cell or in liquid media are known to the art.

25 Another aspect of the invention provides for the positive selection of cells that include a replication competent polynucleotide. In this aspect of the invention, a replication competent polynucleotide typically includes a second coding sequence encoding a selectable marker, and the cell which includes the replication competent polynucleotide is incubated in the presence of a selecting
30 agent. Those cells that can replicate in the presence of the selecting agent contain a polynucleotide that is replication competent. The cells that can replicate are detected by allowing resistant cells to grow in the presence of the selecting agent, and observing, for instance, the presence of colonies and/or the expression of a marker, such as SEAP.

In some aspects, the method may further include isolating virus particles from the cells that contain a replication competent polynucleotide and exposing a second cell to the isolated virus particle under conditions such that the virus particle is introduced to the cell. After providing time for expression of the selectable marker, the second cell is then incubated with the selecting agent. The presence of a cell that replicates indicates the replication competent polynucleotide produces infectious virus particles.

In another aspect, the invention provides a method for detecting a replication competent polynucleotide. The method includes incubating a cell that contains a replication competent polynucleotide of the present invention. The polynucleotide may include a second coding region encoding a selectable or detectable marker. Optionally, the polynucleotide may include a transactivator that interacts with the operator sequence present in the cell. In this aspect, the cell may include a transactivated coding region and an operator sequence operably linked to the transactivated coding region. The method further includes detecting the presence of increased amounts of the replication competent polynucleotide, or the presence or absence of the marker encoded by the second coding sequence or the transactivated coding region present in the cell. The presence of increased amounts of the replication competent polynucleotide or the marker indicates the cell includes a replication competent polynucleotide.

The methods described above for identifying a replication competent polynucleotide can also be used for identifying a variant replication competent polynucleotide, i.e., a replication competent polynucleotide that is derived from a replication competent polynucleotide of the present invention. A variant replication competent polynucleotide may have a faster replication rate than the parent or input polynucleotide. The method takes advantage of the inherently high mutation rate of RNA replication. It is expected that during continued culture of a replication competent polynucleotide in cultured cells, the polynucleotide of the present invention may mutate, and some mutations will result in polynucleotides with greater replication rates. The method includes identifying a cell that has greater expression of a polypeptide encoded by a replication competent polynucleotide. A polynucleotide of the present invention that replicates at a faster rate will result in more of the polynucleotide in the cell, or will result in more of the polypeptide(s) that is encoded by the second coding

region present in the polynucleotide. For instance, when a replication competent polynucleotide encodes a selectable marker, a cell containing a variant polynucleotide having a greater replication rate will be resistant to higher levels of an appropriate selecting agent. When a polynucleotide encodes a transactivator, a cell containing a variant polynucleotide having a greater replication rate than the parent or input polynucleotide will express higher amounts of the transactivated coding region that is present in the cell.

A cDNA molecule of a variant replication competent polynucleotide can be cloned using methods known to the art (see, for instance, Yanagi et al., *Proc. Natl. Acad. Sci., USA*, 94, 8738-8743 (1997)). The nucleotide sequence of the cloned cDNA can be determined using methods known to the art, and compared with that of the input RNA. This allows identification of mutations that have occurred in association with passage of the replication competent polynucleotide in cell culture. For example, using methods known to the art, including longrange RT-PCR, extended portions of a variant replication competent polynucleotide genome can be obtained. Multiple clones could be obtained from each segment of the genome, and the dominant sequence present in the culture determined. Mutations that are identified by this approach can then be reintroduced into the background of the cDNA encoding the parent or input polynucleotide.

The present invention also provides methods for identifying a compound that inhibits replication of a replication competent polynucleotide. The method includes contacting a cell containing a replication competent polynucleotide with a compound and incubating the cell under conditions that permit replication of the replication competent polynucleotide in the absence of the compound. After a period of time sufficient to allow replication of the polynucleotide, the replication competent polynucleotide is detected. A decrease in the presence of replication competent polynucleotide in the cell contacted with the compound relative to the presence of replication competent polynucleotide in a cell not contacted by the compound indicates the compound inhibits replication of the polynucleotide. A compound that inhibits replication of such a polynucleotide includes compounds that completely prevent replication, as well as compounds that decrease replication. Preferably, a compound inhibits replication of a replication competent polynucleotide by at least about 50%, more preferably at least about 75%, most preferably at least about 95%.

The compounds added to a cell can be a wide range of molecules and is not a limiting aspect of the invention. Compounds include, for instance, a polyketide, a non-ribosomal peptide, a polypeptide, a polynucleotide (for instance an antisense oligonucleotide or ribozyme), other organic molecules, or a
5 combination thereof. The sources for compounds to be screened can include, for example, chemical compound libraries, fermentation media of *Streptomyces*, other bacteria and fungi, and extracts of eukaryotic or prokaryotic cells. When the compound is added to the cell is also not a limiting aspect of the invention. For instance, the compound can be added to a cell that contains a replication
10 competent polynucleotide. Alternatively, the compound can be added to a cell before or at the same time that the replication competent polynucleotide is introduced to the cell.

Typically, the ability of a compound to inhibit replication of a replication competent polynucleotide is measured using methods described herein. For
15 instance, methods that use nucleic acid amplification to detect the amount of a replication competent polynucleotide in a cell can be used. Alternatively, methods that detect or select for a marker encoded by a replication competent polynucleotide or encoded by a cell containing a replication competent polynucleotide can be used.

20 In some aspects of the invention, the replication competent polynucleotide of the invention can be used to produce viral particles. Preferably, the viral particles are infectious. For instance, a cell that includes a replication competent polynucleotide can be incubated under conditions that allow the polynucleotide to replicate, and the viral particles that are produced can be isolated using methods
25 routine and known to the art. The viral particles can be used as a source of virus particles for various assays, including evaluating methods for inactivating particles, excluding particles from serum, identifying a neutralizing compound, and as an antigen for use in detecting anti-HCV antibodies in an animal. An example of using a viral particle as an antigen includes use as a positive-control in
30 assays that test for the presence of anti-HCV antibodies.

For instance, the activity of compounds that neutralize or inactivate the particles can be evaluated by measuring the ability of the molecule to prevent the particles from infecting cells growing in culture or in cells in an animal. Inactivating compounds include detergents and solvents that solubilize the

envelope of a viral particle. Inactivating compounds are often used in the production of blood products and cell-free blood products. Examples of compounds that can be neutralizing include a polyketide, a non-ribosomal peptide, a polypeptide (for instance, an antibody), a polynucleotide (for instance, an antisense oligonucleotide or ribozyme), or other organic molecules. Preferably, a neutralizing compound is an antibody, including polyclonal and monoclonal antibodies, as well as variations thereof including, for instance, single chain antibodies and Fab fragments.

Viral particles produced by replication competent polynucleotide of the invention can be used to produce antibodies. Laboratory methods for producing polyclonal and monoclonal antibodies are known in the art (see, for instance, Harlow E. et al. *Antibodies: A laboratory manual* Cold Spring Harbor Laboratory Press, Cold Spring Harbor (1988) and Ausubel, R.M., ed. *Current Protocols in Molecular Biology* (1994)), and include, for instance, immunizing an animal with a virus particle. Antibodies produced using the viral particles of the invention can be used to detect the presence of viral particles in biological samples. For instance, the presence of viral particles in blood products and cell-free blood products can be determined using the antibodies.

The present invention further includes methods of treating an animal including administering neutralizing antibodies. The antibodies can be used to prevent infection (prophylactically) or to treat infection (therapeutically), and optionally can be used in conjunction with other molecules used to prevent or treat infection. The neutralizing antibodies can be mixed with pharmaceutically acceptable excipients or carriers. Suitable excipients include but are not limited to water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, neutralizing antibodies and pharmaceutically acceptable excipients or carriers may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance the effectiveness of the neutralizing antibodies. Such additional formulations and modes of administration as are known in the art may also be used.

The virus particles produced by replication competent polynucleotide of the invention can be used as a source of viral antigen to measure the presence and amount of antibody present in an animal. Assays are available that measure the

presence in an animal of antibody directed to HCV, and include, for instance, ELISA assays and recombinant immunoblot assay. These types of assays can be used to detect whether an animal has been exposed to HCV, and/or whether the animal may have an active HCV infection. However, these assays do not use

5 virus particles, but rather individual or multiple viral polypeptides expressed from recombinant cDNA that are not in the form of virus particles. Hence they are generally unable to detect potentially important antibodies directed against surface epitopes of the envelope polypeptides, nor are they typically measures of functionally important viral neutralizing antibodies. Such antibodies are generally

10 detected with the use of infectious virus particles, such as those that are produced in this system. The use of infectious viral particles as antigen in assays that detect the presence of specific antibodies by virtue of their ability to block the infection of cells with HCV viral particles, or that possibly bind to whole virus particles in an ELISA assay or radioimmunoassay, will allow the detection of functionally

15 important viral neutralizing antibodies.

The present invention also provides a kit for identifying a compound that inhibits replication of a replication competent polynucleotide. The kit includes a replication competent polynucleotide as described herein, and a cell that contains a polynucleotide including a transactivated coding sequence encoding a detectable

20 marker and an operator sequence operably linked to the transactivated coding sequence in a suitable packaging material. Optionally, other reagents such as buffers and solutions needed to practice the invention are also included. Instructions for use of the packaged materials are also typically included.

As used herein, the phrase "packaging material" refers to one or more

25 physical structures used to house the contents of the kit. The packaging material is constructed by well known methods, preferably to provide a sterile, contaminant-free environment. The packaging material may include a label which indicates that the replication competent polynucleotide can be used for identifying a compound that inhibits replication of such a polynucleotide. In addition, the

30 packaging material may contain instructions indicating how the materials within the kit are employed. As used herein, the term "package" refers to a solid matrix or material such as glass, plastic, and the like, capable of holding within fixed limits the replication competent virus and the vertebrate cell.

The present invention is illustrated by the following examples. It is to be understood that the particular examples, materials, amounts, and procedures are to be interpreted broadly in accordance with the scope and spirit of the invention as set forth herein.

5

EXAMPLES

Materials and Methods

Cells. Huh7 cells were grown in Dulbecco's modified Eagle's medium (Gibco BRL, Carlsbad, CA) supplemented with 10% fetal calf serum, penicillin and streptomycin. En5-3 is a clonal cell line derived from Huh7 cells by stable transformation with the plasmid pLTR-SEAP (Yi et al., Virology, 304,197-210 (2002)). These cells were cultured in Dulbecco's modified Eagle's medium (Gibco BRL) supplemented with 10% fetal calf serum, 2 µg/ml blasticidin (Invitrogen), penicillin and streptomycin. Cell lines were passaged once or twice per week. G418 at a concentration of 250 µg/ml was used to select colonies from En5-3 cells transfected with replicon RNAs containing 1a sequences.

Plasmids. The plasmid pBpp-Htat2ANeo was constructed by replacing the *BsrGI-XbaI* fragment of pBpp-Ntat2ANeo/SI (identical to Ntat2ANeo/SI as described by Yi et al. (Yi et al., Virology, 304,197-210 (2002)) with the analogous segment of pH77c (GenBank AF011751) (Yanagi et al., Proc Natl Acad Sci USA, 94, 8738-43 (1997)) engineered to contain a *BsrGI* site at the corresponding location by Quick-Change (Stratagene, La Jolla, CA) mutagenesis. This fragment swap results in the NS3-NS5B sequence in pBpp-Htat2ANeo being identical to that of pH77c, with the exception of the RNA encoding the N-terminal 75 amino acid residues of NS3 that retains the genotype 1b Con1 sequence. Since Bpp-Ntat2ANeo/SI was originally engineered to contain the genotype 1a 5' nontranslated RNA (5'NTR) sequence (Yi et al., Virology, 304,197-210 (2002)), the resulting pBpp-Htat2ANeo construct possesses both a genotype 1a 5'NTR and 1a 3'NTR sequence. Overlapping PCR was used to fuse an anti-genomic hepatitis delta ribozyme sequence directly to the 3' end of the genotype 1a 3'NTR, in order to generate a self-cleaving 3' sequence with the exact 3' terminal nucleotide of HCV (Perrotta and Been, Nucleic Acids Res, 24,1314-21 (1996)). Derivatives of

pBpp-Htat2ANeo containing the adaptive mutations K1691R or S2204I were created by Quick-Change (Stratagene) mutagenesis.

- To construct pBpp-H34A-Ntat2ANeo/SI, an *EcoRI* restriction site was created in pBpp-Ntat2ANeo/SI near the 3' end of the NS4A coding region by Quick-Change mutagenesis. After digestion of the resulting plasmid with *BsrGI* and *EcoRI*, the excised HCV segment was replaced with the equivalent sequence from pH77c which had been amplified by PCR using primers pairs containing terminal *BsrGI* and *EcoRI* sites, respectively. To construct the plasmid Hpp-H34A-Ntat2ANeo, DNA fragments representing the encephalomyocarditis virus (EMCV) internal ribosome entry site (IRES) sequence and the genotype 1a H77c NS3 protein-coding sequence were fused by overlapping PCR. The resulting fragment was digested with *KpnI* at a site located within the EMCV IRES and *BsrGI* at the site created within the modified pH77c NS3 region (see above), then inserted in place of the corresponding fragment in pBpp-H34A-Ntat2ANeo/SI.
- The adaptive mutations, Q1067R or G1188R, were introduced into pHpp-H34A-Ntat2ANeo/SI in a similar fashion, using cDNA fragments prepared by RT-PCR of template RNAs isolated from independent G418-resistant replicon cell lines selected after transfection of En5-3 cells with Hpp-H34A-Ntat2ANeo RNA. pHtat2ANeo/SI was constructed by replacing the *BsrGI-XbaI* fragment of pHpp-H34A-Ntat2ANeo/SI with that of pBpp-Htat2ANeo/SI. A similar strategy was used to construct pHtat2ANeo/QR/SI, pHtat2ANeo/KR/SI, and pHtat2ANeo/QR/KR/SI. Quick-Change (Stratagene) mutagenesis was used to introduce the P1496L, F2080V and K2040R mutations into replicon constructs derived from pHtat2ANeo/SI.
- Modified pH77c plasmids containing adaptive mutations were created by replacing the *BsrGI-XbaI* fragment with the corresponding fragment from the pHtat2ANeo plasmid derivative containing the indicated mutation, except for the Q1067R mutation which was introduced by Quick-Change (Stratagene) mutagenesis. Each mutation was confirmed by sequence analysis. For use as controls, replication-incompetent subgenomic and genome-length genotype 1a constructs (Htat2ANeo/QR/VI/KR/KR5A/SI/AAG and H77/QR/VI/KR/KR5A/SI/AAG) were created by replacing residues 2737-2739 of NS5B ('GDD') with 'AAG' using a similar strategy. Each mutation was confirmed by sequence analysis.

RNA transcription and transfection. RNA was synthesized with T7 MEGAScript reagents (Ambion, Austin, TX), after linearizing plasmids with *Xba*I. Following treatment with RNase-free DNase to remove template DNA and precipitation of the RNA with lithium chloride, the RNA was transfected into

5 Huh7 cells or En5-3 cells by electroporation. Briefly, 5 µg RNA was mixed with 2×10^6 cells suspended in 500 µl phosphate buffered saline, in a cuvette with a gap width of 0.2 cm (Bio-Rad). Electroporation was with two pulses of current delivered by the Gene Pulser II electroporation device (Bio-Rad), set at 1.5 kV, 25 µF, and maximum resistance. For transient replication assays, no G418 was added

10 to the media. Transfected cells were transferred to two wells of a 6-well tissue culture plate, and culture medium removed completely every 24 hrs and saved at 4°C for subsequent SEAP assay. The cells were washed twice with PBS prior to re-feeding with fresh culture medium. Since the culture medium was replaced every 24 hours in these transient assays, the SEAP activity measured in these

15 fluids reflected the daily production of SEAP by the cells. Cells were split 5 days after transfection. Samples of media were stored at 4°C until assayed for SEAP activity at the conclusion of the experiment.

Alkaline phosphatase assay. SEAP activity was measured in 10 µl aliquots of transfected cell supernatant culture fluids using the Phospha-Light

20 Chemiluminescent Reporter Assay (Applied Biosystems/Tropix, Foster City, CA) with the manufacturer's suggested protocol reduced in scale. The luminescent signal was read using a TD-20/20 Luminometer (Turner Designs, Inc., Sunnyvale, CA).

Sequence analysis of cDNA from replicating HCV RNAs. HCV RNA was

25 extracted from cells, converted to cDNA and amplified by PCR as described previously (Yi et al., J Virol, 77, 57-68 (2003)). First-strand cDNA synthesis was carried out with Superscript II reverse transcriptase (Gibco-BRL); pfu-Turbo DNA polymerase (Stratagene) was used for PCR amplification of the DNA. The amplified DNAs were subjected to direct sequencing using an ABI 9600

30 automatic DNA sequencer.

In vitro translation. In vitro transcribed RNA, prepared as described above, was used to program in vitro translation reactions in rabbit reticulocyte lysate (Promega, Madison, WI). Approximately 1 µg RNA, 2 µl of [³⁵S]-methionine (1,000 Ci/mmol at 10 mCi/ml), and 1 µl of an amino acid mixture

lacking methionine were included in each 50 µl reaction mixture. Translation was carried out at 30° C for 90 minutes. Translation products were separated by SDS-PAGE followed by autoradiography or PhosphorImager (Molecular Dynamics) analysis.

5 *Indirect immunofluorescence.* Cells were grown on chamber slides until 70-80% confluent, washed 3 times with PBS, and fixed in methanol/acetone (1:1 V/V) for 10 min at room temperature. A 1:20 dilution of a primary, murine monoclonal antibody to core or NS5A (Maine Biotechnology Services, Portland, ME) was prepared in PBS containing 3% bovine serum albumin, and incubated
10 with the fixed cells for 1 hour at room temperature. Following additional washes with PBS, specific antibody binding was detected with a goat anti-mouse IgG FITC-conjugated secondary antibody (Sigma, St. Louis, Missouri) diluted 1:70. Cells were washed with PBS, counterstained with DAPI, and mounted in Vectashield mounting medium (Vector Laboratories, Burlingame, CA) prior to
15 examination by a Zeiss AxioPlan2 Fluorescence microscope.

Northern analysis for HCV RNA. Replicon-bearing cells were seeded into 10 cm dishes at a density of 5×10^5 cells/dish, and harvested the RNA 4 days later. Total cellular RNA was extracted with Trizol reagent (Gibco-BRL) and quantified by spectrophotometry at 260 nm. Thirty µg of the total RNA extracted from each
20 well was loaded onto a denaturing agarose-formaldehyde gel, subjected to electrophoresis and transferred to positively-charged Hybond-N+ nylon membranes (Amersham-Pharmacia Biotech) using reagents provided with the NorthernMax Kit (Ambion). RNAs were immobilized on the membranes by UV-crosslinking. The membrane was hybridized with a mixture of [32 P]-labeled
25 antisense riboprobe complementary to the 3'-end of the HCV NS5B sequence (nucleotides 8990-9275) derived from pH77C or pHCV-N, and the hybridized probe was detected by exposure to X-ray film.

Results

30 *Transient replication of 1a replicon containing chimeric NS3-coding sequence.* In contrast to genotype 1b HCV, several previous reports suggest that it is difficult to generate subgenomic genotype 1a replicons that are capable of efficient replication in Huh7 cells (Blight et al., Science, 290:1972-4 (2000), Guo et al., J Virol, 75, 8516-23 (2001), Ikeda et al., J Virol, 76, 2997-3006 (2002),

Lanford et al., J Virol, 77,1092-104 (2003)). Similar results were encountered with a dicistronic SEAP reporter replicon constructed from the H77c infectious molecular clone (Yanagi et al., Proc Natl Acad Sci USA, 94, 8738-43 (1997)) that encoded both the HIV tat protein and neomycin phosphotransferase in the upstream cistron. The organization of this latter replicon, Htat2ANeo/SI (Fig. 1), was similar to that of the efficiently replicating, genotype 1b Bpp-Ntat2ANeo/SI replicon (Fig. 1), referred to previously simply as "Ntat2ANeo/SI" (Yi et al., Virology, 304,197-210 (2002)). Most of the HCV polyprotein-coding sequence in Bpp-Ntat2ANeo/SI was derived from the genotype 1b HCV-N strain of HCV (Beard et al., Hepatol., 30, 316-24 (1999)), but the "Bpp" prefix used here and throughout this communication refers to the presence of 225 nucleotides (nts) of sequence that are derived from the Con1 strain of HCV at the extreme 5' end of the polyprotein coding region ("pp" indicates the 5' proximal protease-coding region, Fig. 1). In contrast, all of the HCV sequence in Htat2ANeo/SI (Fig. 1) is derived from the genotype 1a H77c virus, including both the 5' NTR and 3' NTR sequences. Unlike Bpp-Ntat2ANeo/SI RNA, Htat2ANeo/SI RNA did not transduce the selection of G418-resistant colonies, nor induce secretion of SEAP above that observed with a replication-incompetent NS5B-deletion mutant (Δ GDD) when transfected into En5-3 cells (stably transformed Huh7 cells that express SEAP under control of the HIV long terminal repeat promoter) (Yi et al., Virology, 304,197-210 (2002)). in a transient replication assay. This was the case even though the replicon was engineered to contain the genotype 1b adaptive mutation, S2204I, within NS5A (Fig. 1). The absence of apparent replication of Htat2ANeo/SI RNA was striking given the fact that it was derived from a well-documented infectious molecular clone of the H77c strain of HCV (Yanagi et al., Proc Natl Acad Sci USA, 94, 8738-43 (1997)).

Recent reports suggest that the EMCV IRES-driven translation of the second cistron in dicistronic, subgenomic RNAs such as those shown in Fig. 1 may be reduced when the translated RNA sequence is derived from genotype 1a virus, rather than genotype 1b (Gu et al., J Virol, 77, 5352-9 (2003), Guo et al., J Virol, 75, 8516-23 (2001), Lanford et al., J Virol, 77,1092-104 (2003)). However, even when translation of the second cistron is rendered more efficient by replacing the 5' 225 nts of the genotype 1a NS3 sequence with related sequence from the Con1 genotype 1b virus, replication typically has not been observed when the

remainder of the replicon sequence is derived from a genotype 1a virus (Guo et al., J Virol, 75, 8516-23 (2001), Lanford et al., J Virol, 77,1092-104 (2003)). However, Gu et al. (Gu et al., J Virol, 77, 5352-9 (2003)) recently described the successful selection of a replication competent, chimeric replicon in which the 5' 225 nts of the NS3 coding sequence was derived from genotype 1b virus, and the remainder of the second cistron from genotype 1a HCV (construction of chimeric replicons being simplified by a unique *BsrGI* site within the genotype 1b Con1 virus sequence, 225 nts downstream from the 5' end of the NS3 region). This replicon also contained 5'NTR sequence derived from genotype 1b virus, and had a single base change within the genotype 1a 3'NTR sequence. The results of Gu et al. (Gu et al., J Virol, 77, 5352-9 (2003)) suggest that the inclusion of the Con1 sequence at the 5' end of the NS3 region may in some way facilitate replication of the 1a RNA. This hypothesis is strengthened by observations made with genotype 1b replicons derived from HCV-N. Those described previously, including Bpp-Ntat2ANeo/SI RNA, were constructed by ligation of HCV-N sequence to a Con1 replicon at the *BsrGI* site (Guo et al., J Virol, 75, 8516-23 (2001), Ikeda et al., J Virol, 76, 2997-3006 (2002), Yi et al., Virology, 304,197-210 (2002)), and thus they contain 5' proximal NS3 sequence (proximal protease sequence or 'pp', Fig. 1) derived from the Con1 virus. Although this chimeric Con1/HCV-N RNA replicates significantly more efficiently than the originally-described Con1 replicons, the replacement of the 5' proximal NS3 sequence in Bpp-Ntat2ANeo/SI with sequence from HCV-N (resulting in Npp-Ntat2ANeo/SI) virtually ablated its replication phenotype in transient transfection assays, although it remained possible to select G418-resistant colonies at a low frequency following transfection.

To formally assess the ability of the 5' proximal genotype 1b NS3 sequence to enhance genotype 1a RNA replication, the 5' 225 nts of NS3 coding region in Htat2ANeo/SI were replaced with the Con1 sequence, generating Bpp-Htat2ANeo/SI (Fig. 1). The construct was also modified by replacing the *XbaI* restriction site at the 3' end of the HCV sequence with the hepatitis delta virus ribozyme sequence (Perrotta and Been, Nucleic Acids Res, 24,1314-21 (1996)). We have shown previously that the presence of the 4 extraneous nts at the 3' end of the replicon RNA that results from run-off transcription of *XbaI*-digested plasmid DNA reduces the replication competence of genotype 1b RNAs by 2-3

fold (Yi and Lemon, *Rna*, 9, 331-45 (2003)). The inclusion of the ribozyme resulted in self-cleaving RNA transcripts capable of generating the exact 3' terminal HCV RNA sequence. Nonetheless, this modified Bpp-Htat2ANeo/SI RNA still remained incapable of inducing the expression of SEAP in transfected EN5-3 cells beyond that observed following transfection of the Δ GDD RNA. Transfection resulted only in an initial burst in SEAP expression due to translation of the input replicon RNA, without the sustained SEAP expression that is indicative of RNA replication (Fig. 2). However, the Bpp-Htat2ANeo/SI RNA was capable of transducing the selection of G418-resistant cell colonies supporting replication of the RNA over a period of 3-4 weeks following transfection of the cells.

The sequence of replicon RNAs extracted from two independent G418-resistant cell clones selected following the transfection of En5-3 cells with Bpp-Htat2ANeo RNA was analyzed. The presence of a single Lys to Arg mutation located within the NS4A region, at residue 1691 (K1691R) of the polyprotein in both cell clones was determined. This residue is located just beyond the 3' limits of the NS4A cofactor peptide sequence which participates in forming a noncovalent complex with NS3 and enhances its protease activity (Wright-Minogue et al., *J Hepatol*, 32, 497-504 (2000), Yao et al., *Structure Fold Des*, 7, 1353-63 (1999)). To determine whether the K1691R mutation facilitated replication of the chimeric genotype 1b/1a RNA in En5-3 cells, this mutation was introduced into the parental Bpp-Htat2ANeo/SI construct, thereby creating Bpp-Htat2ANeo/KR/SI (see Table 2 for a list of all adaptive mutations identified in these studies, as well as the symbols used to indicate their presence in constructs). As shown in Fig. 2, this single mutation significantly enhanced the replication capacity of the RNA, allowing replication to be detected by a sustained increase in SEAP expression following transient transfection of EN5-3 cells in the absence of G418 (Fig. 2). Since the level of SEAP production has been shown to correlate closely with intracellular replicon RNA abundance in this reporter system (Yi et al., *Virology*, 304, 197-210 (2002), Yi et al., *J Virol*, 77, 57-68 (2003)) we conclude that K1691R is an adaptive mutation. Interestingly, this mutation has been shown previously to confer an enhanced replication phenotype on Con1 replicons (Lohmann et al., *J Virol*, 77, 3007-19 (2003)), whereas the sequence of HCV-N is naturally Arg at this position (Beard et al., *Hepatol.*, 30, 316-24

(1999)). Our results stand in contrast to those reported by Gu et al. (Gu et al., J Virol, 77, 5352-9 (2003)), who identified several mutations within the NS3, NS5A, and NS5B sequences of chimeric genotype 1b-1a RNAs. None of these mutations appeared to enhance the ability of the chimeric RNA to replicate or
5 transduce colony selection.

The 5' 225 nts of the genotype 1a NS3 sequence down modulate replicon amplification. The results described above, as well as those of Gu et al. (Gu et al., J Virol, 77, 5352-9 (2003)) suggest that first 225 nts of the genotype 1a NS3 sequence have a negative impact on the replication of subgenomic HCV replicons.
10 This could occur by down modulation of EMCV IRES-directed translation of the nonstructural proteins (Guo et al., J Virol, 75, 8516-23 (2001)), or by directly influencing replication itself, possibly by influencing an NS3-related function. To address this issue, the identification of additional adaptive mutations capable of compensating for the presence of the 5' proximal genotype 1a protease sequence
15 was sought. Thus additional chimeric replicons containing the entire genotype 1a NS3/4A sequence within the background of Bpp-Ntat2ANeo/SI (Hpp-H34A-Ntat2ANeo/SI, Fig. 3A) were constructed. Also constructed was a variant of this construct in which the first 225 nts of the NS3/4A sequence was replaced with Con1 sequence (Bpp-H34A-Ntat2ANeo/SI, Fig. 3A). In both chimeric RNAs, the
20 sequence extending from NS4B to the 3'NTR was derived entirely from the genotype 1b HCV-N strain. While the replicon containing the entire genotype 1a NS3/4A sequence (Hpp-H34A-NtatNeo/SI) did not show evidence of replication in a transient transfection assay, the variant containing the first 225 nts of the Con1 sequence (Bpp-H34A-NtatNeo/SI) replicated as well as the reference Bpp-
25 Ntat2ANeo/SI replicon (Fig. 3B). This result confirms that the 5' 225 nucleotides of the genotype 1a NS3 sequence have a negative effect on RNA replication in En5-3 cells, and also indicates that the downstream genotype 1a NS3/4A sequence functions well in this context.

Interestingly, despite the lack of detectable RNA replication in the
30 transient assay, selection of stable G418-resistant cell clones following transfection of Hpp-H34A-Ntat2ANeo/SI RNA was possible. Sequencing of replicon RNAs derived from two independent cell clones revealed only a single potentially adaptive mutation in each: Q1067R and G1188R, both of which are located within RNA encoding the NS3 protease (Fig. 3A). The Q1067R mutation

is of particularly interest, since it is within the 5' 225 nucleotides of the NS3 region. When introduced into Hpp-H34A-Ntat2ANeo/SI, both the Q1067R and (to a lesser extent) the G1188R mutations enhanced replication of the RNA to a level that was detectable in the transient assay (Fig. 3B), indicating that both are
5 adaptive mutations and capable of compensating, in part, for the presence of the genotype 1a protease sequence. However, neither of these mutations, when introduced into a replicon containing only genotype 1a sequence (Htat2ANeo/SI), was able to enhance replication to the point where it was evident in the transient assay (Htat2ANeo/QR/SI, Fig. 4).

10 *Transient replication of a genotype 1a replicon in normal Huh7 cells.* To determine whether the K1691R and Q1067R mutations might work cooperatively to confer a transient replication phenotype on the genotype 1a replicon RNA, both were introduced into Htat2ANeo and assessed the ability of the modified RNA to replicate in transfected En5-3 cells. Surprisingly, the combination of the K1691R
15 and Q1067R mutations (in addition to the S2204I mutation in NS5A) conferred a relatively robust replication phenotype on the genotype 1a RNA, such that replication was easily detectable in the transient transfection assay using the SEAP reporter system (Htat2ANeo/QR/KR/SI, Fig. 4B). Using an approach similar to that taken in the preceding experiments, an additional adaptive mutation
20 (F2080V) within the NS5A-coding region (F2080V) was subsequently identified, when cells transfected with Htat2ANeo/QR/KR/SI RNA were subjected to G418 selection pressure. This mutation resulted in slightly greater replication efficiency when introduced into the genotype 1a replicon containing K1691R and Q1067R in addition to S2204I (Htat2ANeo/QR/KR/FV/SI, Fig. 4B). However, F2080V had
25 relatively little effect when added to replicons containing only K1691R or Q1067R (in addition to S2204I) (Fig. 4B). Minimally increased secretion of SEAP above the Δ GDD background was observed during the first 5 days after transfection with Htat2ANeo/KR/FV/SI, but this was no longer apparent after 6 days. The replication phenotype of Htat2ANeo/QR/FV/SI was indistinguishable
30 from that of the replication incompetent Δ GDD mutant in this assay (Fig. 4B). These results are summarized in Fig. 4C.

To facilitate a comparison of these results with those reported previously by Blight et al. (Blight et al., J Virol, 77, 3181-90 (2003)), the adaptive P1496L mutation identified by this group within the helicase domain of NS3 following

transfection of a genotype 1a replicon was introduced into the highly permissive Huh7 subline, Huh-7.5. Consistent with the previous report, a 1a replicon bearing this mutation P1496L demonstrated only minimal evidence of replication in the transient assay (which utilizes En5-3 cells that are comparable to normal Huh7
5 cells in terms of their permissiveness for HCV RNA replication) (Htat2ANeo/PL/SI, Fig. 4B). The addition of the NS5A mutation, F2080V, failed to noticeably enhance the replication capacity of this RNA (Htat2ANeo/PL/FV/SI, Fig. 4B). SEAP expression induced by genotype 1a replicons containing both Q1067R and K1691R was approximately 10-fold that induced by replicons
10 containing P1496L. Since SEAP production from En5-3 cells correlates closely with the intracellular abundance of replicon RNA (Yi et al., Virology, 304,197-210 (2002)), these results suggest that the protease domain mutations make a greater contribution to replication competence of the genotype 1a replicon.

Adaptive mutations within NS3 do not affect EMCV IRES-driven

15 *translation of the second cistron.* As mentioned above, previous reports indicate that the EMCV-driven translation of the second cistron is reduced in genotype 1a replicons in comparison to replicons containing the genotype 1b Con1 sequence (Gu et al., J Virol, 77, 5352-9 (2003), Guo et al., J Virol, 75, 8516-23 (2001), Lanford et al., J Virol, 77,1092-104 (2003)). Although the mechanism is
20 uncertain, the effect appears to be due to the genotype 1a sequence encoding the amino terminus of NS3. Since the adaptive Q1067R mutation is located within this region, we asked whether it or other mutations that enhance 1a replicon amplification do so by improving EMCV IRES-driven translation of the HCV nonstructural proteins. To test this hypothesis, in vitro translation reactions were
25 programmed with genotype 1b and 1a replicon RNAs containing various adaptive mutations, and compared the production of proteins encoded by the second cistron with neomycin phosphotransferase produced from the first cistron. As shown in Fig. 5, the synthesis of NS3 was modestly reduced with replicons containing genotype 1a H77c sequence in the 5' proximal protease region (compare NS3
30 abundance in lanes 4-8 with that in other lanes). However, it was not increased by any of the adaptive mutations, including Q1067R. This result indicates that the difficulty of establishing replication competent 1a replicons is more likely due to the intrinsic property of the 1a sequence, than to an incompatibility of the HCV and EMCV sequences in this region leading to reduced activity of the EMCV

IRES. Nonetheless, the reduced level of translation of the genotype 1a nonstructural proteins that is evident in Fig. 5 may contribute to the poor replication phenotype of these RNAs.

An additional adaptive NS5A mutation further augments replication competence. Although the F2080V mutation in NS5A provided only a slight additional replication advantage to subgenomic genotype 1a RNAs containing the Q1067R, K1691R and S2204I mutations (Fig. 4), additional mutations were subsequently identified concurrently near the C-terminus of NS3 (V1655I) and within NS5A (K2040R) in RNAs replicating within a G418-resistant cell line selected following transfection with the subgenomic Htat2ANeo/QR/KR/SI replicon. As shown in Fig. 6, both of these mutations enhanced the replication capacity of genotype 1a RNA. Addition of the V1655I mutation resulted in a modest enhancement of Htat2ANeo/QR/KR/SI replication, leading to a replication phenotype slightly better than observed with the addition of the F2080V mutation. In contrast, the addition of the K2040R mutation in NS5A resulted in a dramatic increase in replication competence, rendering the replication phenotype of the genotype 1a RNA equivalent to that of the standard genotype 1b HCV-N replicon used in these studies, Bpp-Ntat2ANeo/SI (Fig. 6B). A genotype 1a replicon containing both of these adaptive mutations in addition to those identified earlier replicated with slightly greater efficiency than this reference genotype 1b RNA in the transient assay (Fig. 6B, Htat2ANeo/QR/VI/KR/KR5A/SI). These results were confirmed in independent experiments.

Robust replication of genome-length genotype 1a RNA with adaptive mutations. Encouraged by the above results, we assessed the in vitro replication competence of genomelength, genotype 1a H77c RNA engineered to contain the adaptive mutations described above. As with the dicistronic, subgenomic RNAs, we placed the hepatitis delta ribozyme sequence at the 3' end of the cloned infectious cDNA sequence in pH77c in order to generate RNA transcripts containing an exact HCV 3' terminus. As these genomic RNAs encoded no selectable marker or reporter protein product, their replication was assessed in transfected Huh7 and En5-3 cells by northern blot analysis in comparison with related subgenomic RNAs. Subgenomic and genome-length replication-incompetent H77 mutant RNAs, in which the GDD motif had been replaced with

AAG, served as negative controls for this experiment. For En5-3 cells transfected with the subgenomic RNAs, we also determined levels of SEAP expression.

As expected, the unmodified H77c RNA showed no evidence of replication, even though it has been shown previously to be infectious in chimpanzees when inoculated into liver (Fig. 7, compare lane 7 with the replication defective 1a genomic RNA in lane 11). The introduction of the Q1067R (NS3) mutation, alone or in combination with S2204I (NS5A), was insufficient to confer a detectable level of replication in Huh7 cells. However, when all three mutations were introduced (Q1067R, K1691R and S2204I), the H77c RNA acquired a relatively efficient replication phenotype with readily detectable amplification of the RNA in northern blots of cell lysates prepared 4 days after transfection of either Huh7 or En5-3 cells (Fig. 7, lane 8). Replication of the genome-length RNA was slightly increased by the further addition of the F2080V (NS5A) mutation (Fig. 7, lane 9). However, consistent with the data presented in Fig. 6, the inclusion of both the V1655I mutation in NS3 and the K2040R mutation conferred a substantially more robust replication phenotype on genome-length H77c, when present in combination with other adaptive mutations in NS3, NS4A and NS5A (H77c/QR/VI/KR/KR5A/SI, Fig. 7, compares lane 10 and 11). This experiment thus confirmed the adaptive effects of these mutations. Northern blotting indicated that the replication capacity of genome-length genotype 1a RNAs containing adaptive mutations was significantly greater than the comparable subgenomic, dicistronic genotype 1a replicons, for which the RNA signal 4 days after transfection was low and near the limits of detection in northern blots (Fig. 7, compare lanes 3 to 6 with lanes 8 to 11). These findings are consistent with those reported previously by Blight et al. (*J. Virol.*, 77, 3181-3190 (2003)), and indicate that the inclusion of heterologous sequences in the dicistronic replicons impairs RNA replication competence. Subgenomic replicon RNA was detected unambiguously only in cells transfected with Htat2ANeo/QR/VI/KR/KR5A/SI, the RNA that generated the highest level of SEAP expression (Fig. 7, compare lane 5 and 6).

As a further measure of the replication competence of these modified genome-length H77c RNAs, we also examined transfected En5-3 cells for the presence of core or NS5A proteins using an indirect immunofluorescence method. Introduction of both the K1691R (NS4A) and S2204I mutations resulted in

detectable antigen expression 4 days after transfection, albeit only in a very low percentage of cells (less than 0.01%). However, strong expression of both the core and NS5A proteins was observed in approximately 30% of En5-3 cells 4 days after transfection of RNA containing all four adaptive mutations. Increased
5 replication efficiency of genotype 1a RNAs correlated with a greater proportion of cells supporting the replication of HCV RNA, evidenced by the presence of viral antigen.

Discussion

10 Subgenomic, dicistronic, selectable HCV RNA replicons derived from genotype 1b viruses replicate efficiently in cultured cells (Blight et al, Science, 290:1972-1974 (2000), Guo et al., J. Virol., 75:8516-8523 (2001), Ikeda et al., J. Virol., 76:2997-3006 (2002), Krieger et al., J. Virol., 75:4614-4624 (2001), Lohmann et al., J. Virol., 75:1437-1449 (2001), and Lohmann et al., Science
15 285:110-113 (1999)). These novel RNAs have facilitated the study of HCV RNA replication and substantially accelerated antiviral drug discovery efforts. The Huh7 cell line, derived from a human hepatoma, appears to be uniquely permissive and supportive of the replication of these HCV RNAs, although recent studies suggest that other types of cells may also be permissive for HCV RNA replication
20 (Zhu et al., J. Virol., 77:9204-9210 (2003)). However, despite the success of genotype 1b replicons, it has been difficult to generate RNAs that replicate efficiently in any cell type from other genotypes of HCV, including genotype 1a, (Blight et al, Science, 290:1972-1974 (2000), Guo et al., J. Virol., 75:8516-8523 (2001), Ikeda et al., J. Virol., 76:2997-3006 (2002), and Lanford et al., J. Virol.,
25 77:1092-104 (2003)). This surprising observation indicates that significant biological differences exist between genotype 1a and 1b viruses, despite the fact that the nucleotide sequences of genotype 1a viruses are relatively closely related to those of genotype 1b (~90-93% identity). This biological difference raises the likelihood that antiviral agents that are found to be active against the genotype 1b
30 virus may have significantly lesser activity against genotype 1a viruses. Considering these observations and the relatively high genetic variability that exists between different HCV genotypes, the development of cell culture systems supporting replication of viral RNAs from other genotypes will be important for validating in vitro efficacy of candidate antiviral agents across a range of

genetically distinct HCV genotypes, as well as developing a better overall understanding of these viruses.

Genotype 1a viruses are the most prevalent types of HCV in the United States, and like genotype 1b virus they are relatively refractory to treatment with
5 interferon (Fried et al., *N Engl J Med*, 347, 975-82 (2002), McHutchison and Fried, *Clin Liver Dis*, 7, 149-61 (2003)). Thus far, a detectable level of genotype 1a RNA replication has been reported only in specially isolated, highly permissive Huh7 human hepatoma cell sublines (e.g., Huh-7.5 cells) generated by eliminating the replication of genotype 1b RNA replicons from established replicon cell lines
10 using interferon- α in vitro (Blight et al., *J Virol*, 77, 3181-90 (2003), Grobler et al., *J Biol Chem*, 278,16741-6 (2003)). These previously described genotype 1a RNAs possess cell culture-adaptive mutations that enhance their replication in these special cells, including those selected during the isolation of antibiotic-resistant cell lines containing these 1a replicons (Blight et al., *J Virol*, 77, 3181-90
15 (2003), Grobler et al., *J Biol Chem*, 278,16741-6 (2003)). However, the published reports suggest that these previously described genotype 1a RNAs do not replicate to a detectable level in standard Huh7 cells, and that their capacity for replication in cultured cells is thus limited. In contrast, genotype 1a HCV RNAs are reported here that replicate in a highly efficient manner in normal Huh7 cells.

20 Our results suggest that the highly efficient replication of genotype 1a RNAs requires at least three adaptive mutations located within the NS3, NS4A and NS5A proteins. It is evident that these mutations are mutually reinforcing in their ability to enhance the replication of the genotype 1a RNAs, even though they were identified individually under different circumstances. It was found that the
25 introduction of the S2204I mutation in NS5A, which is known to promote the replication of genotype 1b virus RNAs in Huh7 cells (Blight et al., *Science*, 290:1972-4 (2000)), was not sufficient for subgenomic replicons composed entirely of the genotype 1a sequence to initiate replication in Huh7 cells. However, it made possible the selection of G418-resistant cell colonies following
30 transfection of a chimeric replicon RNA, in which sequence from the infectious molecular clone of the genotype 1a H77c virus encoded all of the nonstructural proteins other than the N-terminal 75 amino acid residues of NS3 which were derived from the genotype 1b Con1 sequence (Fig. 1, Bpp-Htat2ANeo/SI). The HCV RNAs replicating in these cells contained a single mutation within the

NS4A-coding region (K1691R) that enhanced the replication capacity of the original chimeric replicon RNA (Fig. 2). These results suggest that a restriction to the replication of genotype 1a virus in Huh7 cells may reside within the serine protease domain of NS3, since substitution of the N-terminal domain of the genotype 1a protease with that from the Con1 genotype 1b virus allowed the initiation of replication and the selection of G418-resistant cells. A similar conclusion can be drawn from the results reported by Gu et. al. (Gu et al., J Virol, 77, 5352-9 (2003)). Thus, it is interesting that the adaptive mutation K1691R resides within NS4A very close to the surface of the NS3/4A protease complex that it helps to form (Fig. 8).

In an effort to better understand this restriction, a second chimeric replicon containing the complete genotype 1a NS3/4A sequence within the background of a genotype 1b replicon was constructed. This RNA (Hpp-H34A-Ntat2ANeo/SI) did not undergo detectable replication in the transient transfection system utilized in these studies (Fig. 3). However, it was capable of transducing the selection of G418-resistant cell colonies following transfection and antibiotic selection. Analysis of the sequence of the HCV RNAs replicating within these cells identified a second, cell culture-adaptive mutation within the N-terminal region of the NS3 protease (Q1067R), providing further evidence that a primary restriction to replication of genotype 1a virus resides within this domain. Yet additional evidence for this comes from the replication phenotype of the Bpp-H34A-Ntat2ANeo/SI replicon, which also contains all of the genotype 1a NS3/4A sequence except for the N-terminal 75 amino acid residues, and which demonstrated a robust replication phenotype in the transient transfection assay. Thus there appears to be no restriction to replication deriving from inclusion of the genotype 1a NS3 helicase domain, nor for that matter any part of the protease domain except for its N-terminus.

Further work demonstrated that the K1691R and Q1067R mutations worked cooperatively: neither by itself was capable of conferring the capacity for efficient replication on a replicon composed entirely of genotype 1a sequence, but a combination of the two (in addition to the genotype 1b S2204I adaptive mutation) resulted in RNA replication that could be readily detected in the transient transfection assay (Fig. 4). That these mutations should act cooperatively in their effects on replication, as indicated by the data shown in Fig. 4, is

consistent with their location in the polyprotein, since the NS4A protease cofactor domain interacts primarily with residues within the N-terminal domain of the NS3 protease (Wright-Minogue et al., J Hepatol, 32, 497-504 (2000), Yao et al., Structure Fold Des, 7, 1353-63 (1999)).

5 Additional adaptive mutations were identified and verified through an iterative series of experiments involving RNA transfection, isolation of G418-resistant cells, and analysis of the sequence of efficiently replicating genotype 1a RNAs. Also demonstrated was that the S2204I mutation did indeed facilitate the replication of the genotype 1a RNA, as its removal from the efficiently replicating
10 subgenomic RNAs substantially reduced their replication competence in the transient transfection assay. The genotype 1a adaptive mutations identified herein are summarized in Table 2. They can be grouped functionally into two groups: K2040R, F2080V, and S2204I, which are all located within NS5A (a common site of genotype 1b adaptive mutations), and Q1067R, G1188R, V1655I, and K1691R,
15 which are all located in or otherwise associated with the protease domain of NS3. While to some extent solvent exposed, both G1188R and Q1067R are close to the active site of the protease (Fig. 8), and would both add a significant charge difference to the active face of the protein. V1655I is particularly interesting. It is located near the extreme C-terminus of the NS3 protein, downstream of the
20 helicase domain, and close to the protease active site in the crystal structure of the NS3/4A complex (Yao et al., Structure Fold Des, 7, 1353-63 (1999)). In the P3 position of the NS3/4A cleavage site, V1655 is certain to play a role in substrate recognition during the *cis*-active cleavage of the polyprotein at the NS3/4A junction and it remains within the substrate-binding pocket in the crystal structure.
25 The potential impact of the K1691R mutation, within NS4A, on the conformation of the protease active site is much less certain, but it is in close proximity to the NS4A cofactor domain, as mentioned above, and intercalation of this domain into the NS3 protease is well known to modulate the activity of the protease.

Significantly, all of these NS3 and NS4A mutations are located at some
30 distance from other genotype 1a adaptive mutations in NS3 that have been described in the literature (see Fig. 8). These mutations, located at S1222, A1226 and P1496, are all within the helicase domain of NS3 (Blight et al., J Virol, 77, 3181-90 (2003), Grobler et al., J Biol Chem, 278,16741-6 (2003)). While on the surface of the protein, they are located on the side opposite the solvent exposed

surfaces containing the G1188, V1655, and Q1067 residues (Fig. 8). Thus, it is possible that they facilitate genotype 1a RNA replication by a different mechanism than those mutations that cluster near the active site of the protease. At least the P1496L mutation identified by both Blight et al. (Blight et al., J Virol, 77, 3181-90 (2003)) and Grobler et al. (Grobler et al., J Biol Chem, 278,16741-6 (2003)) appears to be substantially less active in conferring replication capacity on the genotype 1a H77c RNA. This was demonstrated by the lack of detectable replication of RNA replicons containing this mutation (Htat2ANeo/PL/SI and Htat2ANeo/PL/FV/SI) in the transient transfection experiment summarized in Fig. 4.

What role could mutations near the active site of the NS3 protease play in promoting the replication of genotype 1a HCV RNA in Huh7 cells? It is unlikely that these mutations work by enhancing translation of the nonstructural proteins under control of the EMCV IRES in the context of the subgenomic replicon, since we observed no difference in translation of these proteins in vitro in reticulocyte lysates programmed with these RNAs (Fig. 5). More importantly, they enhance the replication of genomic H77c RNA lacking any heterologous sequence in Huh7 cells (see Fig. 7). These mutations do not seem likely to promote replication by favorably influencing the ability of the protease to process the viral polyprotein, since the polyprotein segment expressed in the Htat2ANeo derivatives is derived entirely from the same H77c genome, and this replicates very efficiently in chimpanzee liver. However, this does remain a formal possibility that needs to be excluded in future studies. It is possible, instead, that these mutations promote interactions of the NS3/4A complex with specific cellular proteins that play a role in assembly of the viral replicase complex, or otherwise influence replication by disabling innate cellular antiviral defenses.

Foy et al. (Foy et al., Science, 300, 1145-8 (2003)) recently demonstrated that expression of the NS3/4A protease effectively blocked activation of interferon regulatory factor 3 (IRF3) in Huh7 cells infected with Sendai virus, thereby preventing the induction of synthesis of interferon- β and other antiviral cytokines. This immuno-evasive action of NS3 was reversed by a specific ketoamide inhibitor of the NS3/4A protease, and was dependent upon the protease activity of NS3/4A, indicating that NS3/4A is likely to cleave a cellular protein involved in IRF3 signaling following viral infection. While Foy et al. (Foy et al., Science,

300, 1145-8 (2003)) demonstrated that both genotype 1a and genotype 1b proteases are capable of blocking IRF3 activation, it is intriguing to consider that the adaptive mutations within NS3/4A may promote its ability to direct such a cleavage, thereby enhancing replication of the virus by lessening cellular antiviral defenses.

The second group of adaptive mutations identified within NS5A, K2040R, F2080V, and S2204I (Table 2), are likely to function in a fashion similar to NS5A adaptive mutations identified in genotype 1b replicons, which include S2204I. Although their specific mechanism of action is not known, they may either promote the ability of NS5A to assemble a functional replicase complex in Huh7 cells, or perhaps augment the immunomodulatory actions that have been proposed for this viral protein through its interactions with double-stranded RNA stimulated protein kinase R (PKR) (Gale et al., Clin Diagn Virol, 10,157-62 (1998)). The contribution of these adaptive mutations to the replication of the genotype 1a RNA in these studies appears to be additive to that of the NS3/4A mutations (Figs. 3 and 6), not synergistic as shown for the combination of Q1067R and K1691R (Fig. 3).

The complete disclosure of all patents, patent applications, and publications, and electronically available material (including, for instance, nucleotide sequence submissions in, e.g., GenBank and RefSeq, and amino acid sequence submissions in, e.g., SwissProt, PIR, PRF, PDB, and translations from annotated coding regions in GenBank and RefSeq) cited herein are incorporated by reference. The foregoing detailed description and examples have been given for clarity of understanding only. No unnecessary limitations are to be understood therefrom. The invention is not limited to the exact details shown and described, for variations obvious to one skilled in the art will be included within the invention defined by the claims.

All headings are for the convenience of the reader and should not be used to limit the meaning of the text that follows the heading, unless so specified.

Sequence Listing Free Text

- SEQ ID NO:1 Nucleotide sequence of Hepatitis C virus strain H77
- SEQ ID NO:2 Amino acid sequence of HCV polyprotein encoded by nucleotides 342 - 9377 of SEQ ID NO:1.
- 5 SEQ ID NO:3 Nucleotide sequence of Hepatitis C virus strain H
- SEQ ID NO:4 Amino acid sequence of HCV polyprotein encoded by nucleotides 342 - 9377 of SEQ ID NO:3.
- SEQ ID NO:5 HIV tat polypeptide
- SEQ ID NO:6 NS3 recognition site
- 10 SEQ ID NO:7 Nucleotide sequence of HIV SEAP, HIV long terminal repeat (LTR) is depicted at nucleotides 1-719, and secretory alkaline phosphatase is encoded by the nucleotides 748-2239.
- SEQ ID NO:8 Nucleotide sequence of a 3' NTR.
- SEQ ID NO:9 Nucleotide sequence of a 5' NTR
- 15 SEQ ID NO:10 HIV tat polypeptide
- SEQ ID NO:11 genomic length hepatitis C virus, genotype 1a
- SEQ ID NO:12 HCV polyprotein encoded by the coding region present in SEQ ID NO:11.
- SEQ ID NO:13 nucleotide sequence of Htat2ANeo
- 20 SEQ ID NO:14 HCV polyprotein encoded by the coding region present in SEQ ID NO:13.

What is claimed is:

1. A replication competent polynucleotide comprising:
a 5' non-translated region (NTR), a 3' NTR, and a first coding sequence present between the 5' NTR and 3' NTR and encoding a hepatitis C virus polyprotein, wherein the polyprotein comprises an isoleucine at about amino acid 2204, and further comprises an adaptive mutation selected from the group of an arginine at about amino acid 1067, an arginine at about amino acid 1691, valine at about amino acid 2080, an isoleucine at about amino acid 1655, an arginine at about amino acid 2040, an arginine at about amino acid 1188, and a combination thereof.
2. The replication competent polynucleotide of claim 1 further comprising a second coding sequence.
3. The replication competent polynucleotide of claim 2 wherein the second coding sequence encodes a marker.
4. The replication competent polynucleotide of claim 2 wherein the second coding sequence encodes a transactivator.
5. The replication competent polynucleotide of claim 1 wherein the 5' NTR, the 3' NTR, and the nucleotide sequence encoding the polyprotein are genotype 1a.
6. The replication competent polynucleotide of claim 1 wherein the hepatitis C virus polyprotein is a subgenomic hepatitis C virus polyprotein.
7. The replication competent polynucleotide of claim 1 wherein the hepatitis C virus polyprotein comprises cleavage products core, E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B.

8. The replication competent polynucleotide of claim 1 further comprising a nucleotide sequence having cis-acting ribozyme activity, wherein the nucleotide sequence is located 3' of the 3' NTR.
9. A replication competent polynucleotide comprising:
a 5' non-translated region (NTR), a 3' NTR, and a first coding sequence present between the 5' NTR and 3' NTR and encoding a hepatitis C virus polyprotein comprising cleavage products core, E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B, wherein the polyprotein comprises an isoleucine at about amino acid 2204, an arginine at about amino acid 1067, and an arginine at about amino acid 1691.
10. The polynucleotide of claim 9 further comprising a second coding sequence.
11. The polynucleotide of claim 10 wherein the second coding sequence encodes a marker .
12. The polynucleotide of claim 10 wherein the second coding sequence encodes a transactivator.
13. The polynucleotide of claim 9 wherein the 5' NTR, polyprotein, and 3' NTR are genotype 1a.
14. A method for making a replication competent polynucleotide comprising:
providing a polynucleotide comprising a 5' NTR, 3' NTR, a first coding sequence present between the 5' NTR and 3' NTR and encoding a hepatitis C virus polyprotein, wherein the polyprotein comprises a serine at about amino acid 2204, a glutamine at about amino acid 1067, a lysine at about amino acid 1691, a phenylalanine at about amino acid 2080, a valine at about amino acid 1655, a lysine at about amino acid 2040, or a glycine at about amino acid 1188 and wherein the 5' NTR, polyprotein, and 3' NTR are genotype 1a; and
altering the coding sequence such that the polyprotein encoded thereby comprises an isoleucine at amino acid 2204, and at least one adaptive mutation

selected from the group consisting of an arginine at about amino acid 1067, an arginine at about amino acid 1691, a valine at about amino acid 2080, an isoleucine at about amino acid 1655, an arginine at about amino acid 2040, an arginine at about amino acid 1188, and a combination thereof.

15. The method of claim 14 wherein the hepatitis C virus polyprotein is a subgenomic hepatitis C virus polyprotein.

16. The method of claim 14 wherein the hepatitis C virus polyprotein comprises cleavage products core, E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B.

17. A replication competent polynucleotide produced by the method of claim 14.

18. A method for identifying a compound that inhibits replication of a replication competent polynucleotide, the method comprising:

contacting a cell comprising a replication competent polynucleotide with a compound, the replication competent polynucleotide comprising 5' NTR, 3' NTR, a first coding sequence present between the 5' NTR and 3' NTR and encoding a hepatitis C virus polyprotein, wherein the hepatitis C virus polyprotein comprises an isoleucine at about amino acid 2204, and further comprises an adaptive mutation selected from the group of an arginine at about amino acid 1067, an arginine at about amino acid 1691, valine at about amino acid 2080, an isoleucine at about amino acid 1655, an arginine at about amino acid 2040, an arginine at about amino acid 1188, and a combination thereof;

incubating the cell under conditions wherein the replication competent polynucleotide replicates in the absence of the compound; and

detecting the replication competent polynucleotide, wherein a decrease of the replication competent HCV polynucleotide in the cell contacted with the compound compared to the replication competent polynucleotide in a cell not contacted with the compound indicates the compound inhibits replication of the replication competent polynucleotide.

19. The method of claim 18 wherein detecting the replication competent polynucleotide comprises nucleic acid amplification.
20. The method of claim 18 wherein the replication competent polynucleotide further comprises a second coding sequence encoding a marker, and wherein detecting the replication competent polynucleotide comprises identifying the marker.
21. The method of claim 18 wherein the replication competent polynucleotide further comprises a second coding sequence encoding a transactivator, wherein the cell comprises a polynucleotide comprising a transactivated coding sequence encoding a detectable marker and an operator sequence operably linked to the transactivated coding sequence, wherein the transactivator interacts with the operator sequence and alters expression of the transactivated coding sequence, and wherein detecting the replication competent polynucleotide in the cell comprises detecting the detectable marker encoded by the transactivated coding sequence.
22. The method of claim 18 wherein the cell is a human hepatoma cell.
23. The method of claim 18 wherein the hepatitis C virus polyprotein is a subgenomic hepatitis C virus polyprotein.
24. The method of claim 18 wherein the hepatitis C virus polyprotein comprises cleavage products core, E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B.
25. The method of claim 18 wherein the 5' NTR, polyprotein, and 3' NTR are genotype 1a.
26. A method for selecting a replication competent polynucleotide, the method comprising:
 - incubating a cell in the presence of a selecting agent, wherein:
 - the cell comprises a polynucleotide comprising a 5' non-translated region (NTR), a 3' NTR, and a first coding sequence present between

the 5' NTR and 3' NTR and encoding a hepatitis C virus polyprotein, and a second coding sequence, wherein the polyprotein comprises an isoleucine at about amino acid 2204, and further comprises an adaptive mutation selected from the group of an arginine at about amino acid 1067, an arginine at about amino acid 1691, valine at about amino acid 2080, an isoleucine at about amino acid 1655, an arginine at about amino acid 2040, an arginine at about amino acid 1188, and a combination thereof;

the second coding sequence encodes a selectable marker conferring resistance to the selecting agent; and

the selecting agent inhibits replication of a cell that does not express the selectable marker; and

detecting a cell that replicates in the presence of the selecting agent, wherein the presence of such a cell indicates the polynucleotide is replication competent.

27. The method of claim 26 wherein the selecting agent is an antibiotic.

28. The method of claim 26 wherein the cell is a human hepatoma cell.

29. The method of claim 26 wherein the cell is a first cell, the method further comprising:

obtaining a virus particle produced by the first cell;

exposing a second cell to the isolated virus particle and incubating the second cell in the presence of the selecting agent; and

detecting a second cell that replicates in the presence of the selecting agent, wherein the presence of such a cell indicates the replication competent polynucleotide in the first cell produces an infectious virus particle.

30. The method of claim 26 wherein the hepatitis C virus polyprotein is a subgenomic hepatitis C virus polyprotein.

31. The method of claim 26 wherein the hepatitis C virus polyprotein comprises cleavage products core, E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B.

32. The method of claim 26 wherein the 5' NTR, polyprotein, and 3' NTR are genotype 1a.

33. A method for detecting a replication competent polynucleotide, the method comprising:

incubating a cell comprising a replication competent polynucleotide, wherein:

the replication competent polynucleotide comprises a 5' non-translated region (NTR), a 3' NTR, and a first coding sequence present between the 5' NTR and 3' NTR and encoding a hepatitis C virus polyprotein, and a second coding sequence encoding a transactivator, wherein the polyprotein comprises an isoleucine at about amino acid 2204, and further comprises an adaptive mutation selected from the group of an arginine at about amino acid 1067, an arginine at about amino acid 1691, valine at about amino acid 2080, an isoleucine at about amino acid 1655, an arginine at about amino acid 2040, an arginine at about amino acid 1188, and a combination thereof;

the cell comprises a transactivated coding region and an operator sequence operably linked to the transactivated coding region; and

the transactivated coding region encodes a detectable marker, wherein the transactivator alters transcription of the transactivated coding region; and

detecting the detectable marker, wherein the presence of the detectable marker indicates the cell comprises a replication competent polynucleotide.

34. The method of claim 33 wherein the hepatitis C virus polyprotein is a subgenomic hepatitis C virus polyprotein.

35. The method of claim 33 wherein the hepatitis C virus polyprotein comprises cleavage products core, E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B.

36. The method of claim 33 wherein the 5' NTR, polyprotein, and 3' NTR are genotype 1a.

37. A kit comprising:

a replication competent polynucleotide comprising a 5' non-translated region (NTR), a 3' NTR, and a first coding sequence present between the 5' NTR and 3' NTR and encoding a hepatitis C virus polyprotein, and a second coding sequence encoding a transactivator, wherein the polyprotein comprises an isoleucine at about amino acid 2204, and further comprises an adaptive mutation selected from the group of an arginine at about amino acid 1067, an arginine at about amino acid 1691, valine at about amino acid 2080, an isoleucine at about amino acid 1655, an arginine at about amino acid 2040, an arginine at about amino acid 1188, and a combination thereof; and




a cell comprising a polynucleotide comprising a transactivated coding sequence encoding a detectable marker and an operator sequence operably linked to the transactivated coding sequence, wherein the transactivator interacts with the operator sequence and alters expression of the transactivated coding sequence.

38. The method of claim 37 wherein the hepatitis C virus polyprotein is a subgenomic hepatitis C virus polyprotein.


39. The method of claim 37 wherein the hepatitis C virus polyprotein comprises cleavage products core, E1, E2, P7, NS2, NS3, NS4A, NS4B, NS5A, and NS5B.

40. The method of claim 37 wherein the 5' NTR, polyprotein, and 3' NTR are genotype 1a.



1/38

*Fig. 1***Bpp-Ntat2ANeo/SI****Htat2ANeo/SI****Bpp-Htat2ANeo/SI**

2/38


Fig. 2

4/38

Fig. 4A*Fig. 4B**Fig. 4C*


SEAP	NS Substitutions				
	3p	3h	4A	5A	5A
-					SI
-			KR		SI
-	QR				SI
+++	QR		KR		SI
-	QR			FV	SI
+			KR	FV	SI
+++	QR		KR	FV	SI
+		PL			SI
+		PL		FV	SI

5/38

Fig. 5

7/38

Fig. 7



8/38




Fig. 8A*Fig. 8B**Fig. 8C*

Fig. 9

1	ACCTGGAAAA	ACATGGAGCA	ATCACAAAGTA	GCAATACACG	AGCTACCAAT	GCTGCTTTGTG	CCTGGCTAGA	AGCACAAGAG	80
81	GAGGAGGAGG	TGGGTTTTCC	AGTCACACCT	CAGGTACCTT	TAAGACCAAT	GACTTACAAG	GCAGCTGTAG	ATCTTAGCCA	160
161	CTTTTAAAAA	GAAAAGGGGG	GACTGGAAGG	GCTAATTAC	TCCCAAAGAA	GACAAGATAT	CCTTGATCTG	TGGATCTACC	240
241	ACACACAAGG	CTACTTCCCT	GATTAGCAGA	ACTACACACC	AGGGCCAGGG	GTCAGATATC	CACTGACCTT	TGGATGGTGC	320
321	TACAAGCTAG	TACCAGTTGA	GCCAGATAAG	ATAGAAGAGG	CCAATAAAGG	AGAGAACACC	AGCTTGTTAC	ACCTTGTGAG	400
401	CCTGCATGGG	ATGGATGACC	CGGAGAGAGA	AGTGTTAGAG	TGGAGGTTTG	ACAGCCGCCT	AGCATTTTAT	CACGTGGCCC	480
481	GAGAGCTGCA	TCCGGAGTAC	TTCAAGAACT	GCTGACATCG	AGCTTGCTAC	AAGGGACTTT	CCGCTGGGGA	CTTTCCAGGG	560
561	AGGCGTGGCC	TGGCGGGGAC	TGGGGAGTGG	CGAGCCCTCA	GATCCTGCAT	ATAAGCAGCT	GCTTTTGGCC	TGTACTIONT	640
641	CTCTCTGGTT	AGACCAGATC	TGAGCCTGGG	AGCTCTCTGG	CTAACTAGGG	AACCCACTGC	TTAAGCCTCA	ATAAGCTTC	720
721	TGCATGCTGC	TGCTGCTGCT	GCTGCTGGGC	CTGAGGCTAC	AGCTCTCCCT	GGGCATCATC	CCAGTTGAGG	AGGAGAACC	800
801	GGACTTCTGG	AACCGCGAGG	CAGCCGAGGC	CCTGGGTGCC	GCCAAAGAAG	TGCAGCCTGC	ACAGACAGCC	GCCAAGAACC	880
881	TCATCATCTT	CCTGGGCGAT	GGGATGGGG	TGCTACCGT	GACAGCTGCC	AGGATCCTAA	AAGGGCAGAA	GAAGGACAAA	960
961	CTGGGGCCTG	AGATACCCCT	GGCCATGGAC	CGCTTCCCAT	ATGTGGCTCT	GTCCAAGACA	TACAATGTAG	ACAAACATGT	1040
1041	GCCAGACAGT	GGAGCCACAG	CCACGGCCTA	CCTGTGCGGG	GTCAAAGGGCA	ACTTCCAGAC	CATTGGCTTG	AGTGCAGCCG	1120
1121	CCCGCTTTAA	CCAGTGCAAC	ACGACACGCG	GCAACGAGGT	CATCTCCGTG	ATGAATCGGG	CCAAGAAAGC	AGGGAAGTCA	1200
1201	GTGGGAGTGG	TAACCACCAC	ACGAGTGCAG	CACGCCCTCG	CAGCCGGGAC	CTACGCCCAC	ACGGTGAACC	GCAACTGGTA	1280
1281	CTCGGACGCC	GACGTGCCCTG	CCTCGGCCCC	CCAGGAGGGG	TGCCAGGACA	TCGCTACGCA	GCTCATCTCC	AACATGGACA	1360
1361	TTGACGTGAT	CCTAGGTGGA	GGCCGAAAGT	ACATGTTTCC	CATGGGAACC	CCAGACCCCTG	AGTACCCAGA	TGACTACAGC	1440
1441	CAAGGTGGGA	CCAGGCTGGA	CGGGAAGAA	CTGGTGCAGG	AATGGCTGGC	GAAGCGCCAG	GGTGCCCCGT	ATGTGTGGAA	1520
1521	CCGCACTGAG	CTCATGCAGG	CTTCCCTGGA	CCCCCTGTG	ACCATCTCA	TGGGTCTCTT	TGAGCCCTGGA	GACATGAAAT	1600
1601	ACGAGATCCA	CCGAGACTCC	ACACTGGACC	CCTCCCTGAT	GGAGATGACA	GAGGCTGCC	TGCGCCTGCT	GAGCAGGAAC	1680
1681	CCCCGCGGCT	TCCTTCCCTT	CGTGGAGGGT	GGTCGCATCG	ACCATGGTCA	TCATGAAAGC	AGGGCTTACC	GGGCACCTGAC	1760
1761	TGAGACGATC	ATGTTGACG	ACGCCATTGA	GAGGGCGGGC	CAGCTCACCA	CGGAGGAGGA	CACGCTGAGC	CTCGTCACTG	1840
1841	CCGACCACTC	CCACGTCTTC	TCCTTCGGAG	GCTACCCCCCT	CGGAGGGAGC	TCCATCTTCG	GGCTGGCCCC	TGGCAAGGCC	1920
1921	CGGGACAGGA	AGGCTTACAC	GGTCTCTCTA	TACGGAAACG	GTCCAGGCTA	TGTGCTCAAG	GACGGCGCCC	GGCCGGATGT	2000
2001	TACCGAGAGC	GAGAGCGGGA	GCCCCGAGTA	TGGGCAGCAG	TCAGCAGTGC	CCCTGGACGA	AGAGACCCAC	GCAGGCGAGG	2080
2081	ACGTGGCGGT	GTTCCGCGGC	GGCCCCGAGG	CGCACCTGGT	TCACGGCGTG	CAGGAGCAGA	CCTTCATAGC	GCACGTCTATG	2160
22161	GCCTTCGCCG	CCTGCCCTGGA	GCCCTACACC	GCCTGCGACC	TGGCGCCCCC	CGCCGGCACC	ACCGACGCCG	CGCACCCGG	2239

11/38

Fig. 11A-1

1 GCCAGCCCC TGATGGGGC GACACTCCAC CATGAATCAC TCCCCTGTGA 50
 101 TGTCGTGCAG CTCACAGGAG CCCCCTCCC GAGAGAGCCA TAGTGGTCTG 100
 201 GATAAACCG CTAATGCCT GGAGATTGG GGTGCCCCC GAAGACTGC 200
 301 GTACTTCCG GTGCCCCGG AGGTCTCGTA GACCTGTCAC CATGAGCTGC 300
 401 GAGCTTCCG TTCCCGGGT GCGGTTCAGT CATTGTTGGA GTTTACTTGT 400
 501 GAGCGGTGCG AACCTCGAGG TAGACGTCAG CCTATCCCCA AGGCACGTCG 500
 601 GCAATGAGG TTGCGGGTGG GCGGATGCG TCCTGTCTCC CCGTGGCTCT 600
 701 TAAGGTATC GATACCCCTTA CGTGCGGCTT CGCCGACCTC ATGGGGTACA 700
 801 GCGGTCCGGG TTCTGGAAGA CCGCGTGAAC TATGCAACAG GAAACCTTCC 800
 901 TGCCCGCTTC AGCTACCAA GTGCGCAATT CCTCGGGCTT TACCATGTG 900
 1001 CCGTGCACACT CCGGGGTGTG TCATATCGAT TCCCTTGGCT TCGCGAGGGT AACGCCCTGA 1000
 1101 CCCACAACGC AGCTTCGACG TCTTACCTTC TCCCTAGGCT TCGCGAGGGT AACGCCCTGA 1100
 1201 TTGGTCAACT GTTACCTTTC TCCCTAGGCT TCCCTAGGCT TCGCGAGGGT AACGCCCTGA 1200
 1301 GGATATGATG ATGAACCTGT CCCCTACGGC AGCGTTGGT TCGCGAGGGT AACGCCCTGA 1300
 1401 GGAGTCTCTG CCGGATATGC GTATTTCTCC TCCCTAGGCT TCGCGAGGGT AACGCCCTGA 1400
 1501 TCACCGGGGG AAATGCCGGC CGCACACGGC TGAATTGCAA TGAAAGCCCTT AACGCCCTGA 1500
 1601 GCACATCAAT AGCAGGCGCT TGAATTGCAA TGAAAGCCCTT AACGCCCTGA 1600
 1701 GAGAGGTTGG CAGCTGCCG AGCCTTACC GATTTTGGCC GATTTTGGCC 1700
 1801 GGCACATACC TCAAGACCT TGTGGAATG TGCCCGCRAA GAGCGTGTGT 1800
 1901 CAGGTCTGGG GCGCTACCT ACAGTGGGG TGCAAAATGT ACAGTGTGT 1900
 2001 TGGATGAAT CAACTGGATT CACCAAGTG TCGGAGGCGC CCCCCTGTGT 2000
 2101 GCAACATCC GGAAGCCACA TACTCTCGT GCGGTCCGG TCCCTGGATT 2100
 2201 TACCATCAAT TACACCATAT TCAAGTCAG GATGTACGTG GAGGCGGTG 2200
 2301 CTGGAAGACA GGAACAGGTC CGAGCTCAG CCGTTGCTGC TGCCACACC 2300
 2401 CCGGCTCAT CCACTCCAC CAGACATG TGACGCTGCA GTACTGTAC 2400
 2501 TCTCCTGTT CTTCTGCTT CAGACGGCG CAGTGTCTCC GCTTGTGGA 2500
 2601 CTCAATGCAG CATCCCTGG CCGGACGCAC GGTCTTGTGT CTTCTCTCGT 2600
 2701 TCTACGCCCT CTACGGGATG TGGCTCTTCC TCCTGCTCCT GCTGGCGTTG 2700
 2801 CGTTGTTCTT GTCGGGTTAA TGGCGTGCAG TCTGTCCCA TATTACAAG 2800
 2901 GAAGCGCAAC TGACGCTGTG GGTCCCGCC CTTCAACGTC GGGGGGGCG 2900
 3001 ACATCACCAA ACTACTCTG GCAATCTCG GACCCCTTG GATCTTCAA 3000
 3101 GATCTGCGG CTAGCGCGGA AGATAGCGG AGGTCAATAC GTGCAATGG 3100
 3201 ACCCTCTTC GAGCTGGGC GCACACGGC CTGCGAGAT TGCCCGTGGC 3200
 3301 GGGCAGTAT CGCCCGGTG GGTGACATCA TCAACGGCTT CACCGCTCTC 3300
 3401 CAAGGGGTGG AGGTGCTGG GGTGACGTC GGCATACGC CAGCAGACA 3400
 3501 CAAGTGGAGG GTAGGTCCA GATCGTCA ACTGCTACC AACCTTCTC 3500
 3601 CGAGGACCAT CGCATCACCC AAGGTCCTG TCATCCAGAT GTATACCAAT 3600
 3701 GACACCTCTG ACCTGGGGT CCTCGACCT TTACCTGGCT ACAGGCTG 3700
 3801 TGCCCGCGG CCAATTCCCTA CTTGAAGGC TCTCGGGGG GTCCGCTGTT 3800
 3901 TGGGATGGC TAAGGGGTG GACTTATCC CTGTGGAGA CTTAGGACA 3900
 4001 CCAGAGCTTC CAGGTGGCCC ACCTGCATG TCCCACCGGC AGCGGTAAGA 4000

12/38

Fig. 11A-2

4101 CTCAACCCCT CTGTTGCTGC AACGGTGGGC TTTGGTGCTT ACATGTCCAA ACATGTCCAA 4200
 4201 CTGGCAGCCC CATCAGGTAC TCCACCTACG GCAAGTTCTT TGCCGACGGC TGCCGACGGC 4300
 4301 CACGGATGCG ACATCCATCT TGGCATCGG CACTGTCTT GACCACGAG GACCACGAG 4400
 4401 TCCGTCACTC TGCCCATCTC TAACATCGAG GAGGTGCTT TGTCACACAC TGTCACACAC 4500
 4501 GGGGAAGACA TCTCATCTTC TGCCACTCAA AGAAGAGTGC CGACGAGCTC CGACGAGCTC 4600
 4601 TCTTGACGTG TCTGTCAATC CGACAGCGG CGATGTTGTC GTCGTGTCGA GTCGTGTCGA 4700
 4701 TGCAACACGT GTGTCACTCA GACAGTCGAT TTCAGCTCTG ACCCTACCTT ACCCTACCTT 4800
 4801 GCCGGGGCAG GACTGGCAGG GAGTGGCAGG GCATCTATAG CATCTATAG CATCTATAG 4900
 4901 CTATGACGCG GGTGTGCTT GGTATGAGCT CACGCCGCGG GAGACTACAG GAGACTACAG 5000
 5001 CATCTTGAAT TTTGGGAGGG CGTCTTACG GGCCTCACTC ATATAGATGC ATATAGATGC 5100
 5101 TAGCGTACCA AGCCACCGTG TCGCTAGGG CTCAAGCCCC TCCCCATCG TCCCCATCG 5200
 5201 GCCAACACCC CTGCTATACA GACTGGCGC TGTTCAAGAT GAAGTCACCC GAAGTCACCC 5300
 5301 GAGGTCTGTA CGAGCACCTG GGTGCTCGT GCGGCGTCTT TGGCTGCTT TGGCTGCTT 5400
 5401 TCTTGTCGGG GAAGCCGGCA ATTATACCTG ACAGGGAGT TCTCTACCA TCTCTACCA 5500
 5501 AGGATGATG CTCGCTGAGC AGTTCAAGCA GAAGGCCCTC GGCCTCCTC GGCCTCCTC 5600
 5601 TGGCAGAAAC TCGAGGTCTT TTGGGCGAAG CACATGTGGA ATTTATCAG ATTTATCAG 5700
 5701 TTGCTTCATT GATGGCTTTT ACAGCTGCTG TCACCGCGC ACTAACCACT ACTAACCACT 5800
 5801 CGCGCCCGCC GGTCCGCTGA GGTCCGCTG GGTGCTGTT GGTGCTGTT GGTGCTGTT 5900
 5901 GGTATGGCG CGGCGTGTG GTGAGCTCTT GTAGCATTCA AGATCATAG AGATCATAG 6000
 6001 TCTCGCTGG AGCCCTTGTA GTGCTGTGG TCTGCGAGC AATACTGCG AATACTGCG 6100
 6101 AGCCTTCGCC TCCCGGGGGA ACCATGTTC CCCCACGAC TACGTGCCG TACGTGCCG 6200
 6201 ACCAGCTCC TGAGGCGACT GCATCAGTGG ATAGCTCGG AGTGATCCAC AGTGATCCAC 6300
 6301 TGCTGAGCGA CTTTAAGACC TTGCTGAAGC CCAAGCTCAT GCCACAATG GCCACAATG 6400
 6401 AGGAGACGGC ATTATGCACA CTGTGGAGT CTGTGGAGT GAGATCCTG GAGATCCTG 6500
 6501 AACATGTGA GTGGACGTT CCCATTAAC GCCTACACA CGGGCCCTG CGGGCCCTG 6600
 6601 CAGAGGAATA CGTGGAGATA AGGCGGTGG GGGACTTCA CTACGTATCG CTACGTATCG 6700
 6701 ATTTTTCACA GAATGGACG GAATGGACG ACACAGGTTT GCGCCCTCT GCGCCCTCT 6800
 6801 TACCCGGTGG GGTGCAATT ACCTGGAG ACCGAACCGG CCGTAGCCGT CCGTAGCCGT 6900
 6901 GGAGAAGGTT GGCAGAGGG TCACCCCTT CTATGGCGG CTCTCGGCT CTCTCGGCT 7000
 7001 CTCCCTGAC GCGAGCTCA TAGAGGCTAA CCTCTGTGG AGGCAGGAGA AGGCAGGAGA 7100
 7101 GACTCCTTCG ATCCGTTGT GGCAGAGGAG GATGAGCGG AGGTCTCCGT AGGTCTCCGT 7200
 7201 TCTGGCGCG GCGGACTAC AACCCCGCG TAGTAGAGC GTGGAAAAA GTGGAAAAA 7300
 7301 GTCCCTCTCT GTGCTCCG CACGAGGAT TACGGGCGAC GGTACGGTG GTCTACCG GTCTACCG 7400
 7401 AGCTCCTCAA CTTCGGCAT CTTCGGCAT ATACGACCA CATCTCTGA CATCTCTGA 7500
 7501 CCATGCCCCC CTTGGAGGG GAGCCTGGG ATCCGGATC CAGCGACGG CAGCGACGG 7600
 7601 CTCATGCT TATTCCTGGA CAGGCGACT CGTCACCCG TCGCTGCGG TCGCTGCGG 7700
 7701 CACAATCTGG TGTATTCAC CACTTCACG AGTGTGCG AGAGGAGAA AGAGGAGAA 7800
 7801 TGCTCAAGGA GGTCAAGCA CGGCGTCAA AAGTGAAGG TAACCTGTA TAACCTGTA 7900
 7901 GTTGGCTAT GGGCAAAAG ACCTCCCTTG CCATGCCAGA AAGCCCTAG AAGCCCTAG 8000
 8001 ATAGACACTA CCATCATGG CCAAGACTG GTTTCTGCG TTGAGCTGA TTGAGCTGA 8100
 8101 TGGCGTGTG CGAGAAGATG GCCCTGTACG ACGTGGTTAG CAAGCTCCCC CAAGCTCCCC 8200
 8201 GGTGAATTC CTCGTGCAAG CGTGGAGATC CCGATGGGGT CCGATGGGGT CCGATGGGGT 8300

13/38

Fig. 11A-3

```

8301 CGTACGGAGG AGGCAATTTA CCAATGTTGT GACCTGGACC CCAAGCCCG 8400
8401 CCAATTCAAG GGGGAAAC TGCGGCTACC GCAGGTGCCG CCGAGCGGC 8500
8501 GGCAGCCTGT CGAGCCGCAG GGCTCCAGGA CTGCACCATG CTCGTGTGTG 8600
8601 GCGAGCCTGA GAGCCTTCAC GGAGGCTATG ACCAGGTACT CCGCCCCCCC 8700
8701 CCTCCAACT GTCAAGTCGCC CACGACGGCG CTGGAAAGAG GGTCTACTAC 8800
8801 AAGACACACT CCAGTCAATT CCTGGCTAGG CAACATAATC ATGTTTGCCC 8900
8901 ATAGCCAGGG ATCAGCTTGA ACAGGCTCTT AACTGTGAGA TCTACGGAGC 9000
9001 ATGGCCTCAG CGCATTTTCA CTCACAGTT ACTCTCCAGG TGAATCAAT 9100
9101 GAGACACCGG GCCCGAGCG TCCGCGCTAG TCCGCGCTAG GCGCTGGACT 9200
9201 CTCAAACTCA CTCCAATAGC CTGGTCTCTG TTTTGCCCTAC TCGAGGGGTA 9300
9301 CCGGGCCCGG CTGGTCTCTG TTTTGCCCTAC TTTTGCCCTAC TCGAGGGGTA 9400
9401 TCTTAAGCCA TTTCTGTTT TTTTGCCCTAC TTTTGCCCTAC TCGAGGGGTA 9500
9501 TGGTGGCTCC ATCTTAGCCC TAGTCACGGC TAGCTGTGAA AGTCCGTGA 9599

```

[illegible]

Fig. 11B-2

[illegible]

16/38

Fig. 11B-3

3042/901 ATT CTT CAA GCC AGT TTG CTT AAA GTC CCC TAC TTC GTG CGC GTT CAA GGC CTT CTC CGG ATC TGC GCG CTA GCG CGG AAG ATA GCC GGA
 I L Q A S L L L K V P Y F V R R L L L R C A L A R K I A G
 3132/931 GGT CAT TAC GTG CAA ATG GCC ATC ATC AAG TTA GGG GCG CTT ACT GGC ACC TAT GTG TAT AAC CAT CTC ACC CCT CTT CGA GAC TGG GCG
 G H Y V Q M A I I K L G A L L G T Y V Y N H L T P L R D W A
 3222/961 CAC AAC GGC CTG CGA GAT CTG GCC GTG GCT GTG GAA CCA GTC GTC TTC TCC CGA ATG GAG ACC AAG CTC ATC ACG TGG GGG GCA GAT ACC
 H N G L R D L A V A V E P V V V F S R M E T K L I T W G A D T
 3312/991 GCC GCG TGC GGT GAC ATC ATC AAC GGC TTG CCC GTC TCT GCC CGT AGG GGC CAG GAG ATA CTG CTT GGG CCA GCC GAC GGA ATG GTC TCC
 A A C G D I I N G L P V S A R R G Q E I L L G P A D G M V S
 3402/1021 AAG GGG TGG AGG TTG CTG GCG CCC ATC ACG GCG TAC GCC CAG CAG ACG AGA GGC CTC CTA GCG TGT ATA ATC ACC AGC CTG ACT GGC CGG
 K G W R L L A P I T A Y A Q Q T R G L L C I I T S L T G R
 3492/1051 GAC AAA AAC CAA GTG GAG GGT GAG GTC CAG ATC GTG TCA ACT GCT ACC CAA ACC TTC CTG GCA ACG TGC ATC AAT GGG GTA TGC TGG ACT
 D K N Q V E G T I I V S T A T Q T F L A T C I N G V C W T
 3582/1081 GTC TAC CAC GGG GCC GGA ACG AGG ACC ATC GCA TCA CCC AAG GGT CCT GTC ATC CAG ATG TAT ACC AAT GTG GAC CAA GAC CTT GTG GGC
 V Y H G A G T R T I A S P K G P V I Q M Y T N V D Q D L V G
 3672/1111 TGG CCC GCT CCT CAA GGT TCC CGC TCA TTG ACA CCC TGT ACC TGC GGC TCC TCG GAC CTT TAC CTG GTC ACG AGG CAC GCC GAT GTC ATT
 W P A P Q G S R S L T P C T C G S D L Y L V T R H A D V I
 3762/1141 CCC GTG CGC CGG CGA GGT GAT AGC AGG GGT AGC CTG CTT TCG CCC CGG CCC ATT TCC TAC TTT AAA GGC TCC TCG GGG GGT CCG CTG TTG
 P V R R G D S R G S L L L S P R P I S Y L K G S S G G P L L
 3852/1171 TGC CCC GCG GGA CAC GCC GTG GGC CTA TTC AGG GCC GCG GTG TGC ACC CGT GGA GTG GCT AAA GCG GTG GAC TTT ATC CCT GTG GAG AAC
 C P A G H A V G L F R A A V C T R G V A K A V D F I P V E N
 3942/1201 CTA GGG ACA ACC ATG AGA TCC CCG GTG TCC CGG ATC TCC CCA CCA GCA GTG CCC CAG AGC TTC CAG GTG GCC CAC CAT CTG CAT GCT
 L G T T M R S P V F T D N S S P P A V P Q S F Q V A H L A
 4032/1231 CCC ACC GGC AGC GGT AAG AGC ACC AAG GTC CCG GCT GCG TAC GCA GCC CAG GGC TAC AAG GTG TTG GTG CTC AAC CCC TCT GTT GCT GCA
 P T G S G K S T K V P A A A Y A Q G Y K V L V L V L N P S V A A
 4122/1261 ACG CTG GGC TTT GGT GCT TAC ATG TCC AAG GCC CAT GGG GTT GAT CCT AAT ATC AGG ACC GGG GTG AGA ACA ATT ACC ACT GGC AGC CCC
 T L G F G A Y M S K A H G V D P N I R T I T T G S P
 4212/1291 ATC ACC TAC GGC AAG TTC CTT GGC GAG TGC TCA GGA GGT GCT TAT GAC ATA ATA ATT TGT GAC GAG TGC CAC TCC
 I T Y S T Y G K F L A D G G C G G G TGC TCA S G G A Y D I I I C D E C H S
 4302/1321 ACG GAT GCC ACA TCC ATC TTG GGC ATC GGC ACT GTC CTT GAC CAA GCA GAG ACT GCG GGG GCG AGA CTG GTT GTG CTC GGC ACT GCT ACC
 T D A T S I L G I G T V L D Q A E T A G A R L V V L A T A T A

Fig. 11B-4

[illegible]

Fig. 11B-5

5742/1801 CTA ACC ACT GGC CAA ACC CTC CTC TTC AAC ATA TTG GGG GGG TGG GTG GCT GCC CAG CTC GGC GCC CCC GGT GCC GCT ACT GCT ACT GCC TTT GTG
 L T T T G Q T T L L F N I L G G W V A A Q L A P A A T A F V
 5832/1831 GGT GCT GGC CTA GCT GGC GCC GCC ATC GGC AGC GTT GGA CTG GGG AAG GTC CTC GTG GAC ATT CTT GCA GGG TAT GGC GCG GGC GTG GCG
 G A G L A G A A I G S V G L G K V L V D I L A G Y G A G V A
 5922/1861 GGA GCT CTT GTA GCA TTC AAG ATC ATG AGC GGT GAG GTC CCC TCC ACC GAG GAC CTG GTC AAT CTG CTG CCC GGC ATC CTC TCG CCT GGA
 G A L V A F K I M S G E V P S T E D L V N L L P A I L S P G
 6012/1891 GCC CTT GTA GTC GGT GTG GTC TGC GCA GCA ATA CTG CGC CGG CAC GTT GGC CCG GGC GAG GGG GCA GTG CAA TGG ATG AAC CGG CTA ATA
 A L V V G G V C A A I L R R H V G P G G G A V Q TGG ATG AAC CGG CTA ATA
 6102/1921 GCC TTC GCC TCC CGG GGG AAC CAT GTT TCC CCC ACG CAC TAC GTG CCG GAG AGC GAT GCA GCC GCC CGC GTC ACT GCC ATA CTC AGC AGC
 A F A S R G G N H V S P T H Y V P E S D A A R V T A I L S
 6192/1951 CTC ACT GTA ACC CAG CTC CTG AGG CGA CTG CAT CAG TGG ATA AGC TCG GAG TGT ACC ACT TGC TCC GGT TCC TGG CTA AGG GAC ATC
 L T V T Q L L R R L H Q W I S S E C T T P C S G S W L R D I
 6282/1981 TGG GAC TGG ATA TGC GAG GTG CTG AGC GAC TTT AAG ACC TGG CTG AAA GCC AAG CTC ATG CCA CAA CTG CCT GGG ATT CCC TTT GTG TCC
 W D W I C E V L S D F K T W L K A GCC A K L M P Q L P G I P F V S
 6372/2011 TGC CAG CGC GGG TAT AGG GGG GTC TGG CGA GGA GAC GGC ATT ATG CAC ACT CGC TGC CAC TGT GGA GCT GAG ATC ACT GGA CAT GTC AAA
 C Q R G Y R G V W R G D G I M H T R C H C G A E I T G G H V K
 6462/2041 AAC GGG ACG ATG AGG ATC GTC GGT CCT AGG ACC TGC AGG AAC ATG TGG AGT GGG ACG TTC CCC ATT AAC GCC TAC ACC ACG GGC CCC TGT
 N G T M R I V G P R T C R N M W S G T F P I N A Y T T G G C P C
 6552/2071 ACT CCC CTT CCT GCG CCG AAC TAT AAG TTC GCG CTG TGG AGG GTG TCT GCA GAG GAA TAC GTG GAG ATA AGG CGG GTG GGG GAC TTC CAC
 T P L P A P N Y K F A L W R V S A E Y V E I R R V G D F H
 6642/2101 TAC GTA TCG GGT ATG ACT ACT GAC AAT CTT AAA TGC CCG TGC CAG ATC CCA TCG CCC GAA TTT TTC ACA GAA TTG GAC GGG GTG CGC CTA
 Y V S G M T T D N L K C P C Q I P S P E F F T E L D G V L D G V R L
 6732/2131 CAC AGG TTT GCG CCC CCT TGC AAG CCC TTG CTG CCG GAG GAT TCA TTC AGA GTA GGA GGA CTC CAC GAG TAC CCG GTG GGG GTG TCG CAA TTA
 H R F A P P C K P L L R E E V S F R V G L H E Y P V G G G GTG GGG AGA AGG TTG
 6822/2161 CCT TGC GAG CCC GAA CCG GAC GTA GCC GTG TTG ACG TCC ATG CTC ACT GAT CCC TCC CAT ATA ACA GCA GAG GCG GCC GGG AGA AGG TTG
 P C E P E P D V A V L T S M L T D P S H I T A E A A G R L
 6912/2191 GCG AGA GGG TCA CCC CCT TCT ATG GCC AGC TCC TCG GCT AGC CAG CTG TCC GCT CCA TCT CTC AAG GCA ACT TGC ACC GCC AAC CAT GAC
 A R G S P P S M A S S A Q L S A P S L K A T C T A N H D
 7002/2221 TCC CCT GAC GCC GAG CTC ATA GAG GCT AAC CTC CTG TGG AGG CAG GAG ATG GGC GGC AAC ATC ACC AGG GTT GAG TCA GAG AAC AAA GTG
 S P D A E L I E A N L L W R Q E M G G N I T R V E S E N K V

Fig. 11B-6

[illegible]

20/38

Fig. 11B-7

8442/2701 GCG AGC GGC GTA CTG ACA ACT AGC TGT GGT AAC ACC CTC ACT TGC TAC ATC AAG GCC CGG GCA GCC TGT CGA GCC GCA GGG CTC CAG GAC
 A S G V L T T S C G N T L T C Y I K A R A A C A A G A A G L Q D
 8532/2731 TGC ACC ATG CTC GTG TGT GGC GAC GAC TTA GTC GTT ATC TGT GAA AGT GCG GGG GTC CAG GAG GAC GCG GCG AGC CTG AGA GCC TTC ACG
 C T M L V C G D L T V V I C E S A G V Q E D A A S L R A F T
 8622/2761 GAG GCT ATG ACC AGG TAC TCC GCC CCC CCC GAG GAC CCC CCA CAA CCA GAA TAC GAC TTG GAG CTT ATA ACA TCA TGC TCC TCC AAC GTG
 E A M T R Y S A P P G D P P Q P E Y D L E L I T S C S S N V
 8712/2791 TCA GTC GCC CAC GAC GGC GCT GGA AAG AGG GTC TAC TAC CTT ACC CGT GAC CCT ACA ACC CCC CTC GCG AGA GCC GCG TGG TGG GAG ACA GCA
 S V A H D G A G R V Y Y L T T R D P T T T P L A R A A W E T A
 8802/2821 AGA CAC ACT CCA GTC AAT TCC TGG CTA GGC AAC ATA ATC ATG TTT GCC CCC ACA CTG TGG TGG AGG ATG ATA CTG ATG ACC CAT TTC TTT
 R H T P V N S W L G N I I I M F A P T L W A R M I L M T H F F
 8892/2851 AGC GTC CTC ATA GCC AGG GAT CAG CTT GAA CAG GCT CTT AAC TGT GAG ATC TAC GGA GCC GGT TGC TAC TCC ATA GAA CCA CTG GAT CTA CCT
 S V L I A R D Q L E Q A L N C E I Y G A CCA CTC AAT AGG GTG GCC GCA TGC CTC AGA
 8982/2881 CCA ATC ATT CAA AGA CTC CAT GGC CTC AGC GCA TTT TCA CTC CAC AGT TAC TCT CCA GGT GAA ATC AAT AGG GTG GCC GCA TGC CTC AGA
 P I I Q R L H G L S A F S L H S Y S P G E I N R V A A C L E
 9072/2911 GTC CCG CCC TTG CGA GCT TGG AGA CAC CGG GCC CGG AGC GTC CGC GCT AGG CTT CTG TCC AGA GGA GGC AGG GCT GCC ATA
 K L G V P P L R A W R H R A R S R L S R G CTT CTG TCC AGA GGA GGC AGG GCT GCC ATA
 9162/2941 TGT GGC AAG TAC CTC TTC AAC TGG GCA GTA AGA ACA AAG CTC AAA CTC ACT CCA ATA GCG GCT GCG CGG CTG GAC TTG TCC GGT TGG
 C G K Y L F N W A V R T K L K L T P I A A G R L D L S G W
 9252/2971 TTC ACG GCT GGC TAC AGC GGG GGA GAC ATT TAT CAC AGC GTG TCT CAT GCC CGG CCC CGC TGG TTC TGG TTT TGC CTA CTC CTG CTC GCT
 F T A G Y S G G D I Y H S V S H A R P R W F W F C L L L L A
 9342/3001 GCA GGC GTA GGC ATC TAC CTC CTC CCC AAC CGA TGA
 A G V G I Y L L P N R *

21/38

Fig. 12A-1

1	GCAGGCCCC	TGATGGGGC	GACACTCCAC	CATGAATCAC	TCCCTGTGA	GGAACCTACTG	TCTTACGGCA	70	AAAAGCCTTA	GGCATCTCCT	ATGGCAGGAA	GAAGCGGAGA	500
101	TGTCGTCAG	CCTCAGGAC	CCCCCTCC	GGGAGAGCCA	TAGTGTCTG	CGGAACCGGT	GAGTACACCG	80	GAATGGCCAG	GAATGGCCAG	GACGACCGGG	TTCTCCGGC	700
201	GATAAACCCG	CTCAATGCC	GGGATTTGG	CGGTGCCCC	GCAAGACTGC	TAGCCGAGTA	GTGTGGGT	90	GCGAAAGGCC	TTGTGGTACT	GCTGTATAGG	300	
301	GTGCTTGGCA	GTGCCCGGG	AGGTCTGTA	GACCGTGAC	CATGGAGCCA	GTAGATCCTA	GACTAGAGCC	100	CTGGAAGCAT	CCAGGAAGTC	AGCCTAAAC	400	
401	TGCTTGATC	AATTGTATT	GTAAAAAGT	TTGCTTTTCA	TGCCAAGTTT	GTTCATAAC	AAAAGCCTTA	1100	GGCATCTCCT	ATGGCAGGAA	GAAGCGGAGA	500	
501	CAGCGACGAA	GACCTCCTCA	AGGCAGTCA	ACTCATCAAG	TTTCTCTATC	AAAGCAACCC	ACCTCCCAAT	1200	CCCGAGGGGA	CCCGACAGGC	CCGAAGGAAG	600	
601	AATTCGACCT	TCTTAAGCTT	GCGGGAGAC	TCGAGTCCAA	CCCTGGGCCC	GGATCTGTTA	ACATGATTTA	1300	ACAAGATGGA	TTGACGCGAG	GTTCCTCCGC	700	
701	CGCTTGGGTG	GAGAGGCTAT	TCGGCTATGA	CTGGGCACAA	CAGACAATCG	GGTCTCTGA	TGCGCCGCTG	1400	TTCCGGGCTGT	CAGCGCAGGG	GCGCCCGGT	800	
801	CTTTTGTGCA	AGACGACCT	GTCCGGTGCC	CTGAATGAAC	TGCAGGACGA	GGCAGCGCGG	CTATCTGGC	1500	TGGCCACGAC	GGGCGTTCT	TGCGCAGCTG	900	
901	TGCTCGACGT	TGTACTGAA	GCGGGAAGG	ACTGGCTGCT	ATTGGGCGAA	GTGCGGGGGC	AGGATCTCCT	1600	GTCACTCTAC	CTTGCTCCTG	CCGAGAAAT	1000	
1001	ATCCATCATG	GCTGATGCAA	TGCGGCGGCT	GCATACGCTT	GATCCGGCTA	GTGCGGGGGC	CGAGCCGAAC	1700	TGTTCCGCCAG	GCTCAAGGCG	CGCATGCCCG	1200	
1101	CGGATGGAAG	CCGGTCTGT	CGATCAGGAT	GATCTGGAC	AAGAGCATCA	GGGGTCCGG	CCAGCCGAAC	1800	TGTTCCGCCAG	GCTCAAGGCG	CGCATGCCCG	1300	
1201	ACGCGGAGGA	TCTCGTCTGT	ACCCATGGG	ATGCTTGCTT	GCGGAATATC	ATGCTGGAAA	ATGGCCGCTT	1900	GCTGACCGCT	TTCTGATTC	ATCGACTGTG	GCGCGCTGGG	1400
1301	TGTGCGGAC	CGCTATCAGG	ACATAGCGTT	GGCTACCCGT	GATATTGCTG	AAGAGCTTGG	CGGCGAATGG	2000	TTACGGTATC	TTCTCGTGT	TTACGGTATC	1500	
1401	GCGGCTCCCG	ATTGCGAGCG	CATCGCCCTC	TATCGCCCTC	TTGACGAGTT	CTTCTGAGTT	TAAACAGACC	2100	CCCTCTAGCG	CCCTCTAGCG	GGATCAATTC	1600	
1501	CCTTCTCTC	CCTCCOCCC	CCCTAACGTT	ACTGGCCGAA	GCGGCTTGA	ATAAGGCCGG	TGTGCGTTTG	2200	ATTTTCCACC	ATTTTCCACC	ATATTGCCGT	1700	
1601	CTTTTGGCAA	TGTAGGGTC	CGGAACCTG	CCCTTGCTT	CTTGACGAGC	ATTCTTAGGG	GTCTTTCCCC	2300	GGATGCAAG	GGATGCAAG	GTCTGTTGAA	1800	
1701	TGCTCTGAAG	GAAGCAGTC	CCTCTGGAAG	TTCTTGAAGA	CAAAACAACGT	CTGTAGCGAC	CTTTTGACG	2400	CAGCGAACC	CAGCGAACC	CGACAGGTGC	1900	
1801	CTCTGCGGCC	AAAAGCCACG	TGTATAAGAT	ACACCTGCAA	AGGCGGCACA	ACCCAGTGC	CACGTTGTGA	2500	TGTGGAAGA	TGTGGAAGA	GTCAATATGC	2000	
1901	TCTCTCTAAG	CGTATTCAAC	AAGGGCTGA	AGGATGCCCA	GAAGGTACCC	CATTGTATGG	GATCTGATCT	2600	TGACATGCT	TGACATGCT	TTACATGCT	2100	
2001	TTAGTCGAGG	TTAAAAACG	TCTAGGCC	CCGAACACG	GGACGCTGGT	TTTCTTTGA	AAAACAGCAT	2700	CGCCCATCAC	CGCCCATCAC	GGCGTACGCC	2200	
2101	CAGCAGACGA	GAGGCTCTCT	AGGTGTATA	ATCACCAAC	TGCTGAGCCG	GGGCGCGGAA	CAAGTGAAGG	2800	CGCATACCC	CGCATACCC	TCATCCAGAT	2300	
2201	AAAGCTTCC	GGCAACGTC	ATCAATGGG	TATGCTGGAC	TGCTTACCAC	CCGCTCATTT	GACACCTGT	2900	ACCTGCGCT	ACCTGCGCT	TTACCTGGTC	2400	
2301	GTATACCAAT	GTGGACCAAG	ACCTTGTGG	CTGGCCCGCT	CCTCAAGGTT	TAGCTGCTTT	TGCGCCCGGC	3000	CCATTTCCTA	CCATTTCCTA	TCCTCGGGGG	2500	
2401	ACGAGGCACG	CCGATGTAT	TCCCGTGCG	CGGCGAGGT	ATAGCAGGGG	GTGTGACCTT	GTGGAGTGGC	3100	TAAAGCGGTA	TAAAGCGGTA	CTGTGGAGAA	2600	
2501	GTCCGCTGTT	GTCCCGCGG	GGACACGCG	TGGGCTTAT	CAGGGCCGCG	CAGCATGCCC	CCAGAGCTTC	3200	CAGGTGGCCC	CAGGTGGCCC	TCCCACCGGC	2700	
2601	CCTAGGGACA	ACCATAAGT	CCCCGGTGT	CACGGACAAC	TCTCTCCAC	GGTGTGGTG	CTCAACCCCT	3300	AACGCTGGC	AACGCTGGC	TTTGGTGGT	2800	
2701	AGCGGTAAAG	GCACCAAGT	CCCCGGTGT	TACGCAGCCC	AGGCTACAA	ACAAATTACCA	CTGGCAGCCC	3400	TCCACCTACG	TCCACCTACG	GCAAGTTCT	2900	
2801	ACATGTCCAA	GGCCCATGG	GTTGATCCTA	ATATCAGGAC	CGGGTGAGA	AGTGCCACTC	CACGGATGCC	3500	ACATCCATCT	ACATCCATCT	TGGGCATCGG	3000	
2901	TGCCGACGGC	GGGTGCTCAG	GAGGTGCTTA	TGACATAATA	ATTGTGACG	CCCTCCGGGG	TCCGTAAGTC	3600	TGTCCTATCC	TGTCCTATCC	GAGGTGCTG	3100	
3001	GACCAAGCAG	AGACTGCGG	GGCGAGACTG	GTGTGCTCG	CCACTGTCTAC	GTGTGACCTT	GGGGAAGACA	3700	TCTCATCTTC	TCTCATCTTC	AGAAGAAGTC	3200	
3101	TGTCACACCA	CGGAGAGAT	CCCTTTTACG	GCAAGGCTAT	CCCCCTCGAG	GTGTGACCTT	GGGGAAGACA	3800	TCTGTATCC	TCTGTATCC	CGATGTTGTC	3300	
3201	CGACGAGCTC	GCCGCGAAG	TGGTCGCTAT	GGGCATCAAT	GCGGTGGCTT	AGTGTGAGT	TGCAACACGT	3900	GACTGGCAGG	GACTGGCAGG	GGGAAGCCAG	GCATCTATAG	3400
3301	GTCGTGTGCA	CCGATGCTCT	CATGACTGGC	TTTACCGGGG	ACTTCGACTC	AGGACTCAAC	GCCGGGGCAG	4000	GGGTGTGCT	GGGTGTGCT	GGGATGAGCT	CACGCCCGCC	3500
3401	ACCTTACCTT	TACCATTTAG	ACAACCCAGC	TCCCCCAGGA	TGCTGTCTCC	TCTGTGAGTG	CTATGACGGG	4100	TTTGGGAGG	TTTGGGAGG	GGTCTTACG	GGCCTCACTC	3600
3501	ATTGTGGGCA	CCGGGGGAGC	GCCCTCCGG	CATGTTGAC	TGCTGTCTCC	TCCTGTGAGT	CTATGACGGG	4200	AGCCACCGTG	AGCCACCGTG	TGCGCTAGGG	CTCAAGCCCC	3700
3601	GAGACTACAG	TAGGCTTACG	AGGCTACATG	AACACCCCGG	GGCTTCCCTG	TGTCAGGAGC	CATCTTGAAT	4300	CTGTATATA	CTGTATATA	GACTGGGCGC	TGTTCAAGAT	3800
3701	ATATAGATAG	CCACTTTTAA	TCCCAGACAA	AGCAGGCTGG	GGAAGACTTT	CCCTTACCTG	TAGCTGTACA	4400	CGAGCACTCT	CGAGCACTCT	GGTGTCTGTT	GCGGCGCTCC	3900
3801	TCCCCTCATG	TGGGACCCAG	TGTGGAAGTG	TTTGTATCCG	CTTAAACCCA	CCCTTACCTG	GCCGAGCTCT	4500	GAAGCCCGGA	GAAGCCCGGA	ATTATACCTG	ACAGGGAGGT	4000
3901	GAAGTCACCC	TGACGCACCC	AATCACCAAA	TACATCATGA	CATGCTATGC	GGCCGACCTG	GAGGTCTGCTA	4600	GAAGCCCGGA	GAAGCCCGGA	ATTATACCTG	ACAGGGAGGT	4100
4001	TGGCTGCTCT	GGCCGCGGAT	TGCTGTGCTA	CAGGCTGCGT	GGTCAATAGT	GGCAGGATCG	TCTTGTCCGG	4700	GAAGCCCGGA	GAAGCCCGGA	ATTATACCTG	ACAGGGAGGT	4200

22/38

Fig. 12A-2

4101	TCTCTACCAG	GAGTTCGATG	AGATGGAAGA	GTGCTCTCAG	CACCTTACCGT	ACATCGAGCA	AGGATGATG	CTCGCTGAGC	CTCGCTGAGC	AGTTCAAGCA	GAAAGGCCCTC	4200
4201	GGCCTCCTGC	AGACCGCGTC	CCGCATGCA	GAGTTATCA	CCCCTGCTGT	CCAGACCAAC	TGGCAGAAAC	TCGAGGTCTT	TTGGGGCGAAG	TTGGGGCGAAG	CACATGTGGA	4300
4301	ATTTTCATCAG	TGGGATACAA	TACTTGGCGG	GCCTGTCAAC	GCTGCCTGGT	AACCCCGCCA	TTGCTTCATT	GATGGCTTTT	ACAGCTGCCG	TCACCAAGCC	TCACCAAGCC	4400
4401	ACTAACCCACT	GGCCAAACCC	TCCTTTTCAA	CATATTGGGG	GGGTGGGTGG	CATCCCTAGT	GGGTGGGTGG	GGTGGGTGGG	GGTGGGTGGG	GGTGGGTGGG	GGTGGGTGGG	4500
4501	CTAGTGGCG	CCGCCATCGG	CAGCGTTGGA	CTGGGGAAGC	TCCTCGTGGG	CTTCTTTCGA	GGGTATGGCG	CGGCGGTGGC	CGGCGGTGGC	GGGAGCTCTT	GTAGCATTTCA	4600
4601	AGATCATGAG	CGGTGAGGTC	CCCTCAGCGG	AGGACCTGGT	CAATCTGTCT	CCCGCATATC	TCTCGCTGGG	AGCCCTGTGA	GTGCGGTGGG	TCTGCGCAGC	TCTGCGCAGC	4700
4701	AATACTGGCG	CGGCACGTTG	GGCCGGGCGA	GGGGGAGTGG	CAATGGATGA	ACCGGCTAAT	AGCCTTCGCC	TCCCGGGGGA	ACCATGTTTC	CCCCACGCAC	CCCCACGCAC	4800
4801	TACGTGCGCG	AGAGCGATGC	AGCCGCCCGC	GTCACTGCCA	TACTCAGCAG	CCTCACTGTA	ACCCAGCTCC	TGAGGGGACT	GCATCAGTGG	ATAAGCTCGG	ATAAGCTCGG	4900
4901	AGTGATACCA	TCCATGCTCC	GGTTCCTGGC	TAAAGGACAT	CTGGGACTGG	ATATGCGAGG	TGCTGAGCGA	CTTTAAGACC	TGGCTGAAAG	CCAAGCTCAT	CCAAGCTCAT	5000
5001	GCACAACTG	CCTGGGATTC	CCTTTGTGTC	CTGCCAGCGC	GGGTATAGGG	GGGTCTGGCG	AGGAGACGGG	ATTTATGCACA	CTCGCTGCCA	CTGTGGAGCT	CTGTGGAGCT	5100
5101	GAGATCACTG	GACATGTCAA	AAACGGGACG	ATGAGGATCG	TCGGTCTCTAG	GACCTGCAGG	AACATGTGGA	GTGGGACGTT	CCCCATTAAAC	GCCTACACCA	GCCTACACCA	5200
5201	CGGGCCCCCTG	TACTCCCCCTT	CCTGCGCGCA	ACTATAAGTT	CGCGCTGTGG	AGGGTGTCTG	CAGAGGAATA	CGTGGAGATA	AGGCGGGTGG	GGGACTTCCA	GGGACTTCCA	5300
5301	CTACGTATCG	GGTATGACTA	CTGACATCTT	TAAATGCCCG	TGCCAGATCC	CATCGCCCGA	ATTTTTCACA	GAATTTGGAG	GGGTGCGCCT	ACACAGGTTT	ACACAGGTTT	5400
5401	GGCCGCCCTT	GCTGCGGAG	GAGGTATCAT	TCAGAGTAGG	TCCAGATAGG	ACTCCACGAG	TACCGGTGGT	GGTCGCAATT	ACCTTGCGAG	CCCGAACCCG	CCCGAACCCG	5500
5501	ACGTAGCCGT	GTTAGCGTCC	ATGCTCATTC	ATCCCTCCCA	TATAACAGCA	GAGCGCGCCG	GGAGAAAGTT	GGCGAGAGGG	TACCCCTTGG	CTATGGCCAG	CTATGGCCAG	5600
5601	CTCCTCGGCT	AGCCAGCTGT	CCGCTCATTC	TCTCAAGGCA	ACTTGCACCG	CCAACCATGA	CTCCCTGAC	GCCGAGCTCA	TAGAGGCTAA	CCTCCTGTGG	CCTCCTGTGG	5700
5701	AGGCAGGAGA	TGGCGGGCAA	CATCACACAG	GTTAGTTCAG	AGAACAAAGT	GGTGATTTCTG	GACTCCTTCG	ATCCGCTTGT	GGCAGAGGAG	GATGAGCGGG	GATGAGCGGG	5800
5801	AGGTCTCCGT	ACCTGCAGAA	ATTTCTGGGA	AGTCTCGGAG	ATTCGCCCGG	GCCTGCCCCG	TCTGGCGCGC	GCCGGACTAC	AACCCGCCCG	TAGTAGAGAC	TAGTAGAGAC	5900
5901	TGGGAAARAG	AATCAACCTT	ATCTACTGCT	TGGCCGAGC	TTGCCACCAA	CACCTTCCAC	AGCTCCTCAA	CTTCCGGGAT	TACGGGCGAC	AATACGACAA	AATACGACAA	6000
6001	GTCTCTACCG	GCCTGCTGCT	CTGCTGCTGC	CCCCCGACTC	CGAGCTGTAG	TCCTATTCTT	CCATGCCCCC	CCTGGAGGGG	GAGCCTGGGG	ATCCGGATCT	ATCCGGATCT	6100
6101	CATCTCTGTA	GCCGCGCCCT	TCTGCTGCTC	CCCCCGACTC	CGAGCTGTAG	TCGTGTGCTG	CTCAATGTCT	TATTCCTGGA	CAGGCGCACT	CGTCACCCCG	CGTCACCCCG	6200
6201	CAGCGACGGG	TCAATGCTGA	CGGTCAAGTGA	TGGGGCCGAC	ACGGAAGATG	GCTACGCCAT	CACAACTCTGG	TGTATTCCAC	CACCTTACCG	AGTGCTTGCC	AGTGCTTGCC	6300
6301	TGCGCTGCGG	AAGAACAAAA	ACTGCCCATC	AACGCACTGA	GCAACTCGTT	TACACGAGAC	TGCTCAAGGA	GGTCAAAAGG	CGGCGTCAA	AACTGAAGGC	AACTGAAGGC	6400
6401	AAAGGCGAGAA	GAAGTTCACA	TTTGACAGAC	TGCAAGTTCT	GGACAGCCAT	CCAAATCCAA	GTTTGGCTAT	GGGGCAAAAG	ACGTCCGTTG	CCATGCCAGA	CCATGCCAGA	6500
6501	TAACTTGCTA	TCCGTAGAGG	AAAGCTTGCAG	AAAGACCTTC	TGGAAGACAG	TGTAACACCA	ATAGACACTA	CCATCATGGC	CAAGAACAGG	GTTTCTTGCG	GTTTCTTGCG	6600
6601	AAGCCGCTGA	GAAGGGGGGT	CGTAAGCCAG	CTCGTCTCAT	CGTGTTCCTC	GACCTGGGCG	TGCGGCTGTG	CGAGAAGATG	GCCCTGTACG	ACGTGGTTAG	ACGTGGTTAG	6700
6701	TTCAGCCTGA	GAAGGGGGGT	CGTAAGCCAG	CTCGTCTCAT	CGTGTTCCTC	CAGGACAGCG	GGTGAATTC	CTCGTGCAAG	CGTGAAGTC	CAAGAAGACC	CAAGAAGACC	6800
6801	CAAGCTCCCC	CTGGCCGCTG	TGGGAGCTC	CTACGGATTG	CAATACCTAC	GAGCGACATC	CGTACGGAGG	AGGCAATTTA	CCAAATGTTG	GACCTGGACC	GACCTGGACC	6900
6901	CGATGGGGT	TCTCGTATGA	TACCCGCTGT	TTTGACTCCA	CAGTCACTGA	GAGCGACATC	CGTACGGAGG	AGGCAATTTA	CCAAATGTTG	GACCTGGACC	GACCTGGACC	7000
7001	CCCAAGCCCC	GTGCGCATC	AACTCCCTCA	CTGAGAGGCT	TTATGTTGGG	GGCCCTCTTA	CCAAATCAAG	GGGGGAAAGG	TGCGGCTACC	GCAGGTGCCG	GCAGGTGCCG	7100
7101	CGGAGCGGCG	GTACTGACAA	CTAGCTGTGG	TAAACCCCTC	ACTTGCTACA	TCAAGGCCCG	GGCAGCTGTG	CGAGCCGCGG	GGCTCCAGGA	CTGCACCATG	CTGCACCATG	7200
7201	CTCGTGTGTG	CGCAGCACTT	AGTCTGTATC	TGTGAAAGTG	CGGGGGTCCA	GGAGACGCG	GGAGGCTGA	GAGCCTTAC	GGAGGCTATG	ACCAGGTACT	ACCAGGTACT	7300
7301	CGCCCCCCCG	CGGGGACCCC	CCCAACACAG	AATACGACTT	GGAGCTTATA	ACATCATGCT	CGGAGCTGA	GAGCCTTAC	GGAGGCTATG	ACCAGGTACT	ACCAGGTACT	7400
7401	GGTCTACTAC	CTTACCCCGT	ACCCCTACAG	CCCCCTCGCG	AGAGCCGCGT	GGAGACACAG	CCTCCACAGT	GTCACTGCGC	CACGACGGCG	CTGGAAGAG	CTGGAAGAG	7500
7501	ATGTTTGGCC	CCACACTGTG	GGCAGGATG	ATACTGATGA	CCCATTTCTT	TAGGCTCTC	ATAGCCAGGG	ATCAGCTTGA	ACAGGCTCTT	AACTGTGAGA	AACTGTGAGA	7600
7601	TCTACCGGAG	CTGCTACTCC	ATAGAACAC	TGGATCTACC	TCCAATCTAT	CAAAAGACTC	ATGGCTCAG	GCATTTTCA	CTCCACAGTT	ACTCTCCAGG	ACTCTCCAGG	7700
7701	TGAATCAAT	AGGTGTGCGG	CATGCCCTAG	AAAACCTTGG	GTCCCGCCCT	TGCGAGCTTG	GAGACACCGG	CCCCGGAGCG	TCCGCGCTAG	GCTTCTGTCC	GCTTCTGTCC	7800
7801	AGAGGAGGCA	GGGCTGCCAT	ATGTGGCAAG	TACCTCTTCA	ACTGGGCAGT	AAGAACAAAG	CTCAAACTCA	CTCCAAATAG	GGCCGCTGGC	CGGCTGGACT	CGGCTGGACT	7900
7901	TGTCCGGTGT	GTTACGCGCT	GGCTACAGG	GGGAGACAT	TTATCACAGC	GTGCTCATG	CCCGGCCCG	CTGCTCTG	TTTTCCTTGG	TTTTCCTTGG	TTTTCCTTGG	8000
8001	TGAGGGGTTA	GGCATCTACC	TCCCTCCCCA	CCGATGAAGG	TTTGGGTAAA	CACCTCCGCC	TCTTAAAGCA	TTTTCCTGTT	TTTTTCTGTT	TTTTTCTGTT	TTTTTCTGTT	8100
8101	TTTTTCTTTT	TTTTTCTTTT	TTTTTCTTTT	TTTTTCTTTT	TTTTTCTTTT	CTTCTTTTAA	TGGTGGGTCC	ATCTTAGCCC	TAGTACGGCG	TAGTACGGCG	TAGTACGGCG	8200
8201	AGGTCCGCTGA	GCCGCAATGAC	TGCAGAGAGT	GCTGATACTG	GCCTCTCTGC	AGATCATGTG	GGTGGGCAATG	GCATCTCCAC	CTCCTCGCGG	CTCCTCGCGG	CTCCTCGCGG	8300

23/38

Fig. 12A-3

8301 GCATCCGAAG GAGGACGCAC GTCCACTCGG ATGGCTAAGG GAGTCTAGAC
8401 GGCCAAATTC GTAATCATGG TCATAGCTGT TTCCCTGTGT AAATTTGTAT
8501 CTGGGGTGCC TAATAGTGA GCTAATCAC ATTAATTGGG TTGGCTCAC
8601 GGCCAAACGC CGGGAGAGG CGGTTTGGGT ATTGGCGCT CTTCGGCTTC
8701 CAGTCACTC AAAGCGGTA ATACGGTAT CCACAGAAT AGGGGATAAC
8801 AAGAGCCGCG TTGTGGCGT TTTTCCATAG CTTCCGCGCC CTTGACGAGC
8901 TAAAGATACC AGGGTTTCC CCTGGAAGC TCCCTCGTGT GCTCTCCTGT
9001 GCGTGGCGCT TTCTCATAGC TCACGCTGTA GGTATCTCAG TTCGGTGTAG
9101 CCGCTGCGCC TTATCCGGTA ACTATCGTCT TGAGTCCAAC CCGTAAGAC
9201 AGGTATGTAG GCGTGTCTAC AGAGTTCTTG AAGTGTGGC CTAACATACGG
9301 CCTTCGGAAA AAGATTGGT AGCTCTTGAT CCGGCAACA AACCACCGCT
9401 AGGATCTCAA GAAGTCTTT TGATCTTTT TACGGGGTCT GACGCTCAGT
9501 ATCTTCACCT AGATCCCTTT AAATTAATAA TGAAGTTTAA AATCAATCTA
9601 AGGCACCTAT CTCAGCGATC TGCTATTTT GTTCATCCAT AGTGCCTGA
9701 CAGTGCTGCA ATGATACCGC GAGACCCACG CTCACCGGCT CCAGATTTAT
9801 ACTTTATCCG CCTCATCCA GTCTATTAAT TGTGCGCGG AAGCTAGAT
9901 GCATCGTGGT GTACGCTCG TCGTTTGGTA TGGCTTCATT CAGCTCCGGT
10001 GGTAGCTCC TTCGGTCTC CGATCGTTGT CAGAAGTAAG TTGGCCGCGAG
10101 CCATCCGTAA GATGCTTTT TGTGACTGCT TAAAGTGTCT CCAAGTCAAT
10201 GGGATAATAC CGCGCACAT AGCAGAACTT TAAAGTGTCT CCAAGTCAAT
10301 CAGTTCGATG TAACCCACTC GTGCACCCAA CTGATCTTCA GCATCTTTTA
10401 AAAAAGGGAA TAAGGCGGAC ACGGAATGT TGAATACTCA TACTCTTCCT
10501 ACATATTTGA ATGATTTTAG AAAATTAAC AAATAGGGGT TCCGCGCAC
10601 GCGGGTGTG GTGGTACGC GCAGCGTGAC CGCTACACTT GCCAGCGCC
10701 GGCTTTCCC GTCAGCTCT AAATCGGGC ATCCCTTTAG GGTCCGATT
10801 CACGTAGTG GCCATCGCCC TGATAGACGG TTTTTCGCCC TTGACGTTG
10901 CAACCTATC TCGGTCTATT CTTTGTATT ATAAGGGATT TTGCCGATT
11001 TTAACAAA TATTAACAAA ATATTACGT TTACAATTTT CCATTCGCCA
11101 ATTACGCCAG CTGGCGAAG GGGGATGTGC TGCAAGGCGA TTAAGTTGGG
11201 AAGTGACTT GGTCAGCGGC CGCTAATACG ACTCACTATA

8400 AGGCCGACTT
8500 AGTGTAAGC
8600 TTAATGAATC
8700 CGAGCGGTAT
8800 GAAACCGTAA
8900 GACACGGATA
9000 CCTTCGGAA
9100 TTCAGCCCCG
9200 TAGCAGAGCG
9300 AAGCCAGTTA
9400 GCAGAAAAAA
9500 ATCAAAAAAG
9600 TTAATCAGTG
9700 CATCTGGCC
9800 TGGTCTGCA
9900 ATTGCTACAG
10000 GCAAAAAAGC
10100 TACTGTCTATG
10200 CGTCAATAC
10300 TGTGAGATC
10400 AAATGCCGCA
10500 ATGAGCGGAT
10600 CATTAGCGC
10700 CACGTTCCGC
10800 GGTGATGGT
10900 GAACAACACT
11000 TAACGCGAAT
11100 CCTCTTCGCT
11200 GGCCAGTGCC

24/38

Fig. 12B-1

2077/1
 ATG GCG CCC ATC ACG GCG TAC GCC CAG CAG ACG AGA GGC CTC CTA GGG TGT ATA ATC ACC AGC CTG ACT GGC CGG GAC AAA AAC CAA GTG
 M A P I T A Y A Q Q T R G L L G C I I T S L T G R D K N Q V
 2167/31
 GAG GGT GAG GTC CAG ATC GTG TCA ACT GCT ACC CAA ACC TTC CTG GCA ACG TGC ATC AAT AAT GGG GTA TGC TGG ACT GTC TAC CAC GGG GGC
 E G Q I V S T A T Q A T F L A T C I N G V C W T V Y H G A
 2257/61
 GGA ACG AGG ACC ATC GCA TCA CCC AAG GGT CCT GTC ATC CAG ATG TAT ACC AAT GTG GAC CAA GAC CTT GTG GGC TGG CCC GCT CCT CAA
 G T I A S P K G P V I Q M Y T N V D Q D L V G W P A P Q
 2347/91
 GGT TCC CGC TCA TTG ACA CCC TGT ACC TGC TCC TCG GGC TCC TCG GAC CTT TAC CTG GTC ACG AGG CAC GCC GAT GTC ATT CCC GTG CGC CGG CGA
 G S R L T P C T C G S S D L Y L V T R H A D V I P V R R
 2437/121
 GGT GAT AGC AGG GGT AGC CTG CTT TCG CCC CCG CCC ATT TCC TAC TTG AAA GGC TCC TCG GGT CCG CTG TTG TGC CCC GCG GGA CAC
 G D S R G S L L S P R P I S Y L K G S S G G P L L C P A G H
 2527/151
 GCC GTG GGC CTA TTC AGG GCC GCG GTG TGC ACC CGT GGA GTG GCT AAA GCG GTG GAC TTT ATC CCT GTG GAG AAC CTA GGG ACA ACC ATG
 A V G L F R A A V C T R G V A K A V D F I P V E N L G T T M
 2617/181
 AGA TCC CCG GTG TTC ACG GAC AAC TCC TCT CCA CCA GCA GTG CCC CAG AGC TTC CAG GTG GCC CAC CTG CAT GCT CCC ACC GGC AGC GGT
 R S P V F T D N S S P A V P Q S F Q V A H L A P T G S G
 2707/211
 AAG AGC ACC AAG GTC CCG GCT GCG TAC GCA GCC CAG GGC TAC AAG GTG TTG GTG CTC AAC CCC TCT GTT GCT GCA ACG CTG GGC TTT GGT
 K S T K V P A A Y A A Q G G Y K V L N P S V A A T L G F G
 2797/241
 GCT TAC ATG TCC AAG GCC CAT GGG GTT GAT CCT AAT ATC AGG ACC GGG GTG AGA ACA ATT ACC ACT GGC AGC CCC ATC ACG TAC TCC ACC
 A Y M S K A H G V D P N I R T G S P I T T G S P I T Y S T
 2887/271
 TAC GGC AAG TTC CTT GCG GAC GGC GGG TGC TCA GGA GGT GCT TAT GAC ATA ATA ATT TGT GAC GAG TGC CAC TCC ACG GAT GCC ACA TCC
 Y G K F L A D G C S G A Y D I I I C D E C H S T D A T S
 2977/301
 ATC TTG GGC ATC GGC ACT GTC CTT GAC CAA GCA GAG ACT GCG GGG GCG AGA CTG GTT GTG CTC GCC ACT GCT ACC CCT CCG GGC TCC GTC
 I L G I G T V L D Q A E T A G A R L V V L A T A T A C C C T P P G S V
 3067/331
 ACT GTG TCC CAT CCT AAC ATC GAG GAG GTT GCT CTG TCC ACC ACC GGA GAG ATC CCC TTT TAC GGC AAG GCT ATC CCC CTC GAG GTG ATC
 T V S H P N I E V A L S T T G E I P F Y G K A I P L E V I
 3157/361
 AAG GGC GGA AGA CAT CTC ATC TTC TGC CAC TCA AAG AAG AAG TGC GAC GAG CTC GCC GCG AAG CTG GTC GCA TTG GGC ATC AAT GCC GTG
 K G G R H L I F C H S K K K C D E L A A K L V A L G I N A V
 3247/391
 GCC TAC TAC CGC GGT CTT GAC GTG TCT GTC ATC CCG ACC AGC GGC GAT GTT GTC GTC GTC GTG TCG ACC GAT GCT CTC ATG ACT GGC TTT ACC
 A Y Y R G L D V S I P T S T D A L M T G F T
 3337/421
 GGC GAC TTC GAC TCT GTG ATA GAC TGC AAC ACG TGT GTC ACT CAG ACA GTC GAT TTC AGC CCT ACC TTT ACC ATT GAG ACA ACC
 G D F D S V I D C N T C V T Q T V D F S L D P T F T I E T T

Fig. 12B-2

3427/451	CAG GAT GCT GTC TCC AGG ACT CAA CGC CGG GGC AGG ACT GGC AGT GGC AGG GGG AAG CCA GGC ATC TAT AGA TTT GTG GCA CCG GGG
3457/461	CAG GAT GCT GTC TCC AGG ACT CAA CGC CGG GGC AGG ACT GGC AGT GGC AGG GGG AAG CCA GGC ATC TAT AGA TTT GTG GCA CCG GGG
3517/481	TCC GGC ATG TTC GAC TCG TCC TCG TCG TCC TGT GAG TGC TAT GAC GCG GGC TGT GCT TGG TAT GAG CTC ACG CCC GCC GAG ACT
3607/511	CGA GCG TAC ATG AAC ACC CCG GGG CTT CCC GTG TGC CAG GAC CAT CTT GAA TTT TGG GAG GGC GTC TTT ACG GGC CTC
3697/541	GAT GCC CAC TTT TTA TCC CAG ACA AAG CAG AGT GGG GAG AAC TTT CCT TAC CTG GTA CCG TAC CAA GCC ACC GTG TGC GCT
3787/571	GCC CCT CCC CCA TCG TGG GAC CAG ATG TGG AAG TGT TTG ATC CGC CTT AAA CCC ACC CTC CAT GGG CCA ACA CCC CTG CTA
3877/601	GCT GGT CAG AAT GAA GTC ACC CTG ACG CAC CCA ATC ACC AAA TAC ATC ATG ACA TGC ATG TCG GCC GAC CTG GAG GTC
3967/631	TGG GTG CTC GTT GGC GGC GTC CTG GCT GCT CTG GCC GCG TAT TGC TGC ATC ACA GGC TGC GTG GTC ATA GTG GGC AGG
4057/661	GGG AAG CCG GCA ATT ATA CCT GAC AGG GAG GTT CTC TAC CAG GAG TTC GAT GAG ATG GAA GAG TGC TCT CAG CAC TTA
4147/691	GGG AAG CCG GCA ATT ATA CCT GAC AGG GAG GTT CTC TAC CAG GAG TTC GAT GAG ATG GAA GAG TGC TCT CAG CAC TTA
4237/721	GCT GTC CAG ACC AAC TGG CAG AAA CTC GAG GTC TTT TGG GCG AAG CAC ATG ATG TGG AAT TTC ATC AGT GGG ATA CAA TAC TTG
4327/751	ACG CTG CCT GGT AAC CCC GGT AAC CCC GGT AAC FTT ACA GCT GCC GTC ACC AGC CCA CTA ACC ACT GGC CAA
4417/781	AAC CTC CTC TTC AAC ATA TTG GGG GGG TGG GTG GCT GCT GCC GGT GGC GCG GCG TAT GGC GCG GGC GTG GGA GCT CTT GTA GCA
4507/811	GGC AGC GTT GGA CTG GGC AAG GTC CTC GTG GAC ATT CTT GCA GGG TAT GGC GCG GGC GTG GGA GCT CTT GTA GCA
4597/841	ATG AGC GGT GAG GTC CCC TCC ACG GAG GAC CTG GTC AAT CTG CTG CCC GGC ATC CTC TCG CCT GGA GCC CTT GTA GTC GGT
4687/871	GCA ATA CTG CGC CGG CAC GTT GGC CCG GGC GAG GGC GCA GTG CAA TGG ATG AAC CGG CTA ATA GCC TTC GCC TCC CGS

Fig. 12B-3

4777/901 GGG AAC CAT GTT TCC CCC ACG CAC TAC GTG CCG GAG AGC GAT GCA GCC GCC GGC GTC ACT GCC ATA CTC AGC AGC CTC ACT GTA ACC CAG
 G N H V S P T H Y V P E S S D A A A R V T A I L S L T T G C C A T A T Q
 4867/931 CTC CTG AGG CGA CTG CAT CAG TGG ATA AGC TCG GAG TGT ACC ACT CCA TGC TCC GGT TCC TGG CTA AGG GAC ATC TGG GAC TGG ATA TGC
 L L R R L H Q W I S S E C T T T P C S G S W L R R D I W D W I C
 4957/961 GAG GTG CTG AGC GAC TTT AAG ACC TGG CTG AAA GCC AAG CTC ATG CCA CAA CTG CCT GGG ATT CCC TTT GTG TCC TGC CAG CGC GGG TAT
 E V L S D F K T W L K A K A K L M P Q L P G I P F V S C Q R G Y
 5047/991 AGG GGG GTC TGG CGA GGA GAC GGC ATT ATG CAC ACT CGC TGC CAC TGT GGA GCT GAG ATC ACT GGA CAT GTC AAA AAC GGG ACG ATG AGG
 R G V W R G D G I M H T R C H C G A E I T G H V K N G T M R
 5137/1021 ATC GTC GGT CCT AGG ACC TGC AGG AAC ATG TGG AGT GGG ACG TTC CCC ATT AAC GCC TAC ACC ACG GGC CCC TGT ACT CCC CTT CCT GCG
 I V G P R T C R N M W S G T F P I N A Y T T G P C T T P L P A
 5227/1051 CCG AAC TAT AAG TTC GCG CTG TGG AGG GTG TCT GCA GAG GAA TAC GTG GAG ATA AGG CCG GTG GGG GAC TTC CAC TAC GTA TCG GGT ATG
 P N Y K F A L W R V S A E Y V E I R R V G D F H Y V S G M
 5317/1081 ACT ACT GAC AAT CTT AAA TGC CCG TGC CAG ATC CCA TCG CCC GAA TTT TTA ACA GAA TTG GAC GGG GTG CGC CTA CAC AGG TTT GCG CCC
 T T D N L K C P C Q I P S P E F T E L D G V R L H R F A P
 5407/1111 CCT TGC AAG CCC TTG CTG CCG GAG GAG GTA TCA TTC AGA GTA GGA CTC CAC GAG TAC CCG GTG GGG TCG CAA TTA CCT TGC GAG CCC GAA
 P C K P L L R E V S F R V G L H E Y P V G S Q L P C E P E
 5497/1141 CCG GAC GTA GCC GTG TTG ACG TCC ATG CTC ACT GAT CCC TCC CAT ATA ACA GCA GAG GCG GCC GGG AGA AGG TTG GCG AGA GGG TCA CCC
 P D V A V L T S M L T D P S H I T A E A G R R L A R G S P
 5587/1171 CCT TCT ATG GCC AGC TCC TCG GCT AGC CAG CTG TCC GCT CCA TCT CTC AAG GCA ACT TGC ACC GCC AAC CAT GAC TCC CCT GAC GCC GAG
 P S M A S S A S Q L S A P S L K A T C T A N H D S P D A E
 5677/1201 CTC ATA GAG GCT AAC CTC CTG TGG AGG CAG GAG ATG GGC GGC AAC ATC ACC AGG GTT GAG TCA GAG AAC AAA GTG GTG ATT CTG GAC TCC
 L I E A N L L W R Q E M G G N I T R V E S E N K V V I L D S
 5767/1231 TTC GAT CCG CTT GTG GCA GAG GAG GAT GAG CCG GAG GTC TCC GTA CCT GCA GAA ATT CTG CGG AAG TCT CGG AGA TTC GCC CGG GCC CTG
 F D P L V A E E D E R E V S V P A E I L R K S R R F A R A L
 5857/1261 CCC GTC TGG GCG CCG GAC TAC AAC CCC CCG CTA GTA GAG ACG TGG AAA AAG CCT GAC TAC GAA CCA CCT GTG GTC CAT GGC TGC CCG
 P V W A R P D Y N P P L V E T W K K P D Y E P P V V H G C P
 5947/1291 CTA CCA CCT CCA CCG TCC CCT CCT GTC CCT CCG CCT CCG AAA AAG CGT ACG GTG GTC CTC ACC GAA TCA ACC CTA TCT ACT GCC TTG GCC
 L P P P R S P P P R K K R T V V L T E S T L S T A L A
 6037/1321 GAG CTT GCC ACC AAA AGT TTT GGC AGC TCC TCA ACT TCC GGC ATT ACG GGC GAC AAT ACG ACA TCC TCT GAG CCC GCC CCT TCT GGC
 E L A T K S F G S S S T S G I T G D N T T S S E P A P S G

Fig. 12B-4

[illegible]

Fig. 12B-5

7477/1801
 AAT TCC TGG CTA GGC AAC ATA ATC ATG TTT GCC CCC ACA CTG TGG GCG AGG ATG ATA CTG ATG ACC CAT TTC TTT AGC GTC CTC ATA GCC
 N S W L G N I I M F A P T L L L M T H F F S V L I A
 7567/1831
 AGG GAT CAG CTT GAA CAG GCT CTT AAC TGT GAG ATC TAC GGA GCC TGC TAC TCC ATA GAA CCA CTG GAT CTA CCT CCA ATC ATT CAA AGA
 R D Q L E Q A L N C E I Y G A C Y S I E P L D L P P I I Q R
 7657/1861
 CTC CAT GGC CTC AGC GCA TTT TCA CTC CAC AGT TAC TCT CCA GGT GAA ATC AAT AGG GTG GCC GCA TGC CTC AGA AAA CTT GGG GTC CCG
 L H G L S A F S L H S Y S P G E I N R V A A C L R K L G V P
 7747/1891
 CCC TTG CGA GCT TGG AGA CAC CGG GCC CGG AGC GTC CGC GCT AGG CTT CTG TCC AGA GGA GGC AGG AAG GGC ATA TGT GGC AAG TAC CTC
 P L R A W R H R A R S V R A R L L S R G G R A A I C G K Y L
 7837/1921
 TTC AAC TGG GCA GTA AGA ACA AAG CTC AAA CTC ACT CCA ATA GCG GCC GCT GGC CGG CTG GAC TTG TCC TCC GGT TGG TTC ACG GCT GGC TAC
 F N W A V R T K L K L T P I A A A G R L D L S G W F T A G Y
 7927/1951
 AGC GGG GGA GAC ATT TAT CAC AGC GTG TCT CAT GCC CGG CCC CGC TGG TTC TGG TTT TGC CTA CTC CTG CTC GCT GCA GGG GTA GGC ATC
 S G G D I Y H S V S H A R P R W F W F C L L L L A A G V G I
 8017/1981
 TAC CTC CTC CCC AAC CGA TGA
 Y L L P N R +

Fig. 13A

29/38

```
1 gccagccccc tgatgggggc gacactccac catgaatcac tcccctgtga ggaactactg
61 tctttcacgca gaaagcgtct agccatggcg ttagtatgag tgtcgtgcag cctccaggac
121 cccccctccc gggagagcca tagtgggtctg cggaaccggt gagtacaccg gaattgccag
181 gacgaccggg tcttttcttg gataaaccog ctcaatgcct ggagatttgg gcgtgcccc
241 gcaagactgc tagccgagta gtgttgggtc gcgaaaggcc ttgtggtact gcctgatagg
301 gtgcttgcca gtgccccggg aggtctcgtg gaccgtgcac catgagcacg aatcctaaac
361 ctcaaagaaa aaccaaacgt aacaccaacc gtgccccaca ggacgtcaag ttcccgggtg
421 gcgctcagat cgttgggtga gtttacttgt tgccgcgcag ggccctaga ttgggtgtgc
481 gcgcgacgag gaagacttcc gagcgggtcg aacctcgagg tagacgtcag cctatcccca
541 aggcacgtcg gcccaggggc aggacctggg ctacgcccgg gtacccttgg cccctctatg
601 gcaatgaggg ttgcgggtgg gcgggatggc tctgtctccc ccgtggctct cggcctagct
661 ggggccccac agacccccgg cgtaggtcgc gcaatttggg taaggctatc gataccctta
721 cgtgcggctt cgccgacctc atggggtaca taccgctcgt cggcggccct ctggaggcg
781 ctgcccaggg cctggcgcat ggcgtccggg ttctggaaga cggcgtgaac tatgcaacag
841 ggaaccttcc tggttgctct ttctctatct tccctctggc cctgctctct tgcctgactg
901 tgcccgcttc agcctacca gtgcgcaatt cctcggggct ttaccatgtc accaatgatt
961 gccctaactc gagtattgtg tacgaggcgg ccgatgccat cctgcacact ccgggggtgtg
1021 tcccttgctg tgcgaggggt aacgcctcga ggtgttgggt ggcggtgacc cccacggtgg
1081 ccaccagggg cggcaaactc cccacaacgc agcttcgacg tcatatcgat ctgcttgtcg
1141 ggagcgccac cctctgctcg gccctctacg tgggggacct gtgcgggtct gtctttcttg
1201 ttggtcaact gtttaccttc tctcccaggc gccactggac gacgcaagac tgcaattgtt
1261 ctatctatcc cggccatata acgggtcctc gcatggcatg ggatatgatg atgaaactgg
1321 cccctacggc agcgttgggt gtagctcagc tgctccggat cccacaagcc atcatggaca
1381 tgatcgctgg tgctcactgg ggagtcctgg cgggcatagc gtatttctcc atgggtggga
1441 actgggcgaa ggtcctggta gtgctgctgc tatttgccgg cgtcgacgcg gaaacccacg
1501 tcaccggggg aaatgccggc cgcaccacgg ctgggcttgt tggctctcct acaccaggcg
1561 ccaagcagaa catccaactg atcaacacca acaccggct ggttagcagg gctcttctat caacacaaat
1621 tgaattgcaa tgaaagcctt aacaccggct ccagctgccg acgccttacc gattttgccc
1681 tcaactcttc aggtgtctct gagaggttgg ccagctgccg cgacgaacgc cctactgct
1741 agggctgggg tctatcagat tatgccacg gaagcggcct gagcgtgtgt ggcccggat
1801 ggcactaccc tccaagacct tgtggcattg tgcccgcaaa gagcgtgtgt ggcccggat
1861 attgcttcac tcccagcccc gtgggtgggt gaacgaccga caggtcgggc gcgcctacct
1921 acagctgggg tgcaaatgat acggatgtct tcgtccttaa caacaccagg ccaccgctgg
1981 gcaattgggt cggttgtacc tggatgaact caactggatt caccaaagtg tgcggagcgc
2041 ccccttgtgt catcgagggg gtgggcaaca acaccttgc ctgccccact gattgcttcc
2101 gcaaacatcc ggaagccaca tactctcggg gcggctccgg tccctggatt acaccagggt
2161 gcatggtcga ctacccgat aggttttggc actatccttg taccatcaat tacaccatat
2221 tcaaagttag gatgtacgtg ggaggggtcg agcacaggct ggaagcggcc tgcaactgga
2281 cgcggggcga acgctgtgat ctggaagaca gggacaggct cgagctcagc ccgttgcctg
2341 tgtccaccac acagtggcag gtccttcctg gttctttcac gacctgcca gccttgcctc
2401 ccggcctcat ccacctccac cagaacattg tggacgtgca gtacttgtag ggggtagggt
2461 caagcatcgc gtcctgggcc attaagtggg agtacgtcgt tctctgttcc cttctgcttg
2521 cagacgcgcg cgtctgctcc tgcttgggga tgatgttact catatcccaa gcggaggcgg
2581 ctttgagaaa cctcgtaata ctcaatgcag catccctggc cgggacgcac ggtcttgtgt
2641 ccttctcgtg gttcttctgc tttgcgtggg atctgaaggg taggtgggtg cccggagcgg
2701 tctacgccct ctacgggatg tggcctctcc tccgtctcct gctggcggtg cctcagcggg
2761 catacgcact ggacacggag gtggccgcgt cgtgtggcgg cgttgttctt gtcgggttaa
2821 tggcgctgac tctgtcgcca tattacaagc gctatatcag ctggtgcatt tgggtgcttc
2881 agtattttct gaccagagta gaagcgcaac tgcacgtgtg ggttcccccc ctcaacgtcc
2941 ggggggggcg cgatgccgtc atcttactca tgtgtgtagt tggttccccg ctggtatttg
3001 acatcaccaa actactcctg gccatcttcg gaccttctg gattcttcaa gccagtttgc
3061 ttaaagtcct ctacttcgtg cgcgttcaag gccttctccg gatctgcgcg ctagcgcgga
```

Fig. 13B

30/38

```

3121 agatagccgg aggtcattac gtgcaaattgg ccatcatcaa gttagggggcg cttactggca
3181 cctatgtgta taaccatctc acccctcttc gagactgggc ctgcagagatc
3241 tggccgtggc tgtggaacca gtcgtcttct cccgaatgga gaccaagctc atcacgtggg
3301 gggcagatac cgccgcgtgc ggtgacatca tcaacggcct gcccgtctct gcccgtaggg
3361 gccaggagat actgcttggg ccagccgacg gaatggtctc caaggggtgg aggttctgtg
3421 cgcccatcac ggcgtacgcc cagcagacga gaggcctcct aggggtgtata atcaccagcc
3481 tgactggccg ggacaaaaac caagtggagg gtgaggtcca gatcgtgtca actgctaccc
3541 aaaccttcct ggcaacgtgc atcaatgggg tatgctggac tgtctaccac ggggccggaa
3601 cgaggaccat cgcatacccc aagggtcctg tcatccagat gtataccaat gtggaccaag
3661 accttgtggg ctggcccgtt cctcaagggt cccgctcatt gacaccctgt acctgcccgt
3721 cctcggacct ttacctggtc acgaggcacg ccgatgtcat tcccgtgcgc cggcgagggtg
3781 atagcagggg tagcctgctt tcgccccggc ccatttccta cttgaaaggc tcctcggggg
3841 gtccgctgtt gtgccccggc ggacacgcgg tgggcctatt cagggccgcg gtgtgcaccc
3901 tgggagtggc taaagcgggt gactttatcc ctgtggagaa cctagggaca accatgagat
3961 ccccggtgtt caccgacaac tcctctccac cagcagtgcc ccagacttc tacgtggccc
4021 acctgcatgc tcccaccggc agcggtaaga gcaccaaggt cccggtgcgc tacgcagccc
4081 agggctacaa ggtgttgggt ctcaaccctt ctgttgctgc aacgctgggc tttggtgctt
4141 acatgtccaa ggcccatggg gttgatccta atatcaggac cggggtgaga acaattacca
4201 ctggcagccc catcacgtac tccacctacg gcaagttcct tgccgacggc ggggtgctcag
4261 gaggtgctta tgacataata atttgtgacg agtgccactc cacggaatgcc acatccatct
4321 tgggcatacg cactgtcctt gaccaagcag agactgcggg ggcgagactg gttgtgctcg
4381 ccaactgtac cctcggggc tcgctactg tgtcccatcc taacatcgag gaggttgcct
4441 tgtccaccac cggagagatc cctttttacg gcaaggctat ccccctcgag gtgatcaagg
4501 ggggaagaca tctcatcttc tgccactcaa agaagaagtg cgacgagctc gccgcgaagc
4561 tggtcgcatt gggcatcaat gccgtggcct actaccgcgg tcttgacgtg tctgtcatcc
4621 cgaccagcgg cgaatgtgtc gtcgtgtcga ccgatgctct catgactggc tttaccggcg
4681 acttcgactc tgtgatagac tgcaacacgt gtgtcactca gacagcttg ttcagccttg
4741 accctacctt taccattgag acaaccacgc tccccaggga tgctgtctcc aggactcaac
4801 gccggggcag gactggcagg ggggaagccag gcatctatag atttgtggca ccgggggagc
4861 gccctccgg catgttcgac tcgtccgtcc tctgtgagtg ctatgacgcg ggctgtgctt
4921 ggtatgagct cagccccgcc gagactacag ttaggctacg agcgtacatg aacaccccg
4981 ggcttccgt gtgccaggac catcttgaat tttgggaggg cgtctttacg ggccctactc
5041 atatagatgc ccacttttta tcccagacaa agcagagtgg ggagaacttt gcttacctgg
5101 tagcgtacca agccaccgtg tgcgttaggg ctcaagcccc tcccctatcg tgggaccaga
5161 tgtggaagtg tttgatccgc cttaaaccce ccctccatgg gccaacaccc ctgctataca
5221 gactgggcgc tgttcagaat gaagtcaccc tgacgcaccc aatcaccaaa tacatcatga
5281 catgcatgtc ggccgacctg gaggtcgtca cgagcacctg ggtgctcgtt ggcggcgtcc
5341 tggctcctc ggccgcgtat tgcctgtcaa caggctgcgt ggtcatagtg ggcaggatcg
5401 tcttgtccgg gaagccggca attatacctg acagggagg tctctaccag gagttcgatg
5461 agatggaaga gtgctctcag cacttacctg acatcgagca agggatgatg ctcgctgagc
5521 agttcaagca gaaggccctc ggccctcctg agaccgcgtc ccgccatgca gaggttatca
5581 cccctgctgt ccagaccaac tggcagaaac tcgaggctct ttgggcgaag cacatgtgga
5641 atttcatcag tgggatacaa tacttggcgg gctgtcaaac gctgcctggt aaccccgcca
5701 ttgcttcatt gatggctttt acagctgcgg tcaccagccc actaaccact ggccaaaccc
5761 tcctcttcaa catattgggg ggggtgggtg ctgcccagct cgccgcccc ctgacctgta
5821 ctgcctttgt ggggtgctggc ctagtggcg cgcctatcgg cagcgttgga ctggggaagg
5881 tcctcgtgga cattcttgca gggataggcg cgggcgtggc gggagctctt gtagcattca
5941 agatcatgag cggtgaggtc ccctccacgg aggacctggt caatctgctg cccgccatcc
6001 tctcgcctgg agcccttgta gtcggtgtgg tctgcgcagc aatactgcgc cggcacgttg
6061 gcccgggcga gggggcagtg caatggatga accggctaata agccttcgcc tcccggggga
6121 accatgttcc cccacgcac cctcactgta agagcgatgc agcgcggcgg gtcactgcca
6181 tactcagcag cctcactgta acccagctcc tggggcact taagggacat gcatcagtgg gtagctcgg
6241 agtgataccac tccatgctcc ggttctctgg taagggacat ctgggactgg atatgcgagg
6301 tgctgagcga ctttaagacc tggctgaaag ccaagctcat gccacaactg cctgggattc
6361 cctttgtgtc ctgccagcgc gggataggg gggctctggc aggagacggc attatgcaca
6421 ctcgctgcca ctgtggagct gagatcactg gacatgtcaa aaacgggacg atgaggatcg
6481 tcggtcctag gacctgcagg aacatgtgga gtgggacgtt ccccataac gcctacacca

```

Fig. 13C

31/38

```

6541 cgggcccctg tactcccctt cctgcgccga actataagtt cgcgctgtgg aggggtgtctg
6601 cagaggaata cgtggagata aggcgggtgg gggacttcca ctacgtatcg ggtatgacta
6661 ctgacaatct taaatgcccg tgccagatcc catcgcccga atttttcaca gaattggacg
6721 ggggtgcgcct acacaggttt gcgccccctt gcaagccctt gctgcgggag gaggtatcat
6781 tcagagtagg actccacgag tacccggtgg ggtcgcaatt accttgcgag cccgaaccgg
6841 acgtagccgt gttgacgtcc atgctcactg atccctccca tataacagca gaggcggccg
6901 ggagaagggt ggcgagaggg tcacccccctt ctatggccag ctccctcggt agccagctgt
6961 ccgctccatc tctcaaggca acttgcaccg ccaacatga ctcccctgac gccgagctca
7021 tagagctaa cctcctgtgg aggcaggaga tgggcggcaa catcaccagg gttgagtcag
7081 agaacaagt ggtgattctg gactccttcg atccgcttgt ggcaaggagg gatgagcggg
7141 aggtctccgt acctgcagaa attctgcgga agtctcgagg attcgccccg gccctgcccc
7201 tctgggcgcg gccggactac aacccccccg tagtagagac gtggaaaaag cctgactacg
7261 aaccacctgt ggtccatggc tgcccgtac cactccacg gtcccctcct gtgctccgc
7321 ctcggaaaaa gcgtacgggtg gtccctcacc aatcaaccct atctactgcc ttggccgagc
7381 ttgccaccaa aagtttttgg agctcctcaa ctccggcat tacgggcgac aatcacgcaa
7441 catcctctga gccgccccct tctggctgcc ccccgcactc cgacgttgag tcctattctt
7501 ccatgcccc cctggagggg gagcctgggg atccggatct cagcgacggg tcatggtcga
7561 cggtcagtag tggggccgac acggaagatg tctgtgtctg ctcaatgtct tattcctgga
7621 caggcgcact cgtcaccccc gtcgctgcgg aagaacaaaa actgccccatc aacgcactga
7681 gcaactcgtt gctacgccat cacaatctgg tgtattccac cacttcacgc agtgcttgcc
7741 aaaggcagaa gaaagtcaca tttgacagac tgcaagttct ggacagccat taccaggacg
7801 tgctcaagga ggtcaaagca gcggcgctcaa aagtgaaggc taacttgcta tccgtagagg
7861 aagcttgtag cctgacgccc ccacattcag ccaaatccaa gtttggtctat ggggcaaaag
7921 acgtccgttg ccatgccaga aaggccgtag ccacatcaa ctccgtgtgg aaagaccttc
7981 tggaagacag tgtaacacca atagacacta ccatcatggc caagaacgag gttttctgcg
8041 ttcagcctga gaaggggggt cgtaagccag ctgctctcat cgtgttcccc gacctgggcg
8101 tgcgcgtagt cgagaagatg gccctgtacg acgtggttag caagctcccc ctggccgtga
8161 tgggaagctc ctacggattc caatactcac caggacagcg ggttgaattc ctgctgcaag
8221 cgtggaagtc caagaagacc ccgatggggg tctcgtagta taccgctgtt tttgactcca
8281 cagtcactga gagcgacatc cgtacggagg aggcaattta ccaatgttgt gacctggacc
8341 cccaagcccc cgtggccatc aagtcctca ctgagaggct ttatgttggg ggccctctta
8401 ccaattcaag gggggaaaaa tgccggtacc gcaggtgccg cgcgagcggc gtactgacaa
8461 ctagctgtgg taacaccctc acttgctaca tcaaggcccc ggagcctgt cgagccgacg
8521 ggtccagga ctgcaccatg ctcggtgtg ggcagcactt agtcgttatc tgtgaaagt
8581 cgggggtcca ggaggacgag ggcagcctga gagccttcac ggaggctatg accaggctat
8641 ccgccccccc cggggacccc ccacaaccag aatacgactt ggagcttata acatcatgct
8701 cctccaacgt gtcagtcgcc cacgacggcg ctggaaagag ggtctactac cttaccctgt
8761 accctacaac cccctcgcg agagccgcgt gggagacagc aagacacact ccagtcaatt
8821 cctggctagg caacataatc atgtttgccc ccacactgtg ggcgaggatg atactgatga
8881 cccatttctt tagcgtcctc atagccagg atcagcttga acaggctctt aactgtgaga
8941 tctacggagc ctgctactcc atagaaccac tggatctacc tccaatcatt caaagactcc
9001 atggcctcag cgcattttca ctccacagtt actctccagg tgaaatcaat aggggtggccg
9061 catgcctcag aaaacttggg gtcccgcctt tgcgagcttg gagacaccgg gcccgagcg
9121 tccgcgctag gcttctgtcc agaggaggca gggctgccat atgtggcaag tacctcttca
9181 actgggcagt aagaacaaag ctcaaactca ctccaatagc ggccgctggc cggtggact
9241 tgtccggtg gttcacggct ggctacagcg gggagacat ttatcacagc gtgtctcatg
9301 cccggccccg ctggttctgg ttttgccctac tctgtctgc tgcaggggtg ggcattctacc
9361 tctcccccaa ccgatgaagg ttggggtaaa cactccggcc tcttaagcca tttcctgttt
9421 tttttttttt tttttttttt tttttctttt ttttttctt tcttttctt ctttttttcc
9481 tttctttttt cttcttttaa tgggtggtcc atcttagccc tagtcacggc tagctgtgaa
9541 aggtccgtga gccgcagac tgcagagagt gctgatactg gcctctctgc agatcatgt

```

Fig. 13D

32/38

MSTNPKPQRKTKRNTNRRPQDVKEPGGGQIVGGVYLLPRRGPRL
GVRATRKTSERSQPRGRRQPIPKARRPEGRTWAQPGYPWPLYGNEGCGWAGWLLSPRG
SRPSWGPTDPRRRSRNLGKVIDTLTCGFADLMGYIPLVGAPLGGAAARALAHGVRVLED
GVNYATGNLPGCSFSIFLLALLSCLTVPASAYQVRNSSGLYHVTNDCPNSSIVYEAAD
AILHTPGCVPCVREGNASRCWVAVTPTVATRDGKLPITTQLRRHIDLLVGSATLCSALY
VGDLGGSVFLVGQLFTFSPRRHWTQDCNCSIYPGHITGHRMAWDMMNWSPTAALV
AQLLRIPQAIMDMIAGAHWGVLAGIAYFSMVGNWAKVLVLLLFAGVDAETHVTGGNA
GRTTAGLVGLLTPGAKQNIQLINTNGSWHINSTALNCNESLNTGWLGLFYQHKNSS
GCPERLASCRRLTDFAGWGPISYANGSGLDERPYCWHYPPRPGIVPAKSVCGPVYC
FTPSPVVVGTTDRSGAPTYSWGANDTDVFLNNTRPPLGNWFGCTWMNSTGFTKVGCA
PPCVIGGVGNNTLLCPTDCFRKHPEATYSRCGSGPWITPRCMVDYPYRLWHYPCTINY
TIFKVRMYVGGVEHRLEAACNWTGERCDLEDNRSELSPLLLSTTQWQVLPSCFTTL
PALSTGLIHLHQNIQNDVQYLYGVGSSIASWAIKWEYVLLFLLADARVCSCLWMMLL
ISQAEAALENLVILNAASLAGTHGLVSFLVFFCFAWYLGKRWVPGAVYALYGMWPLLL
LLLALPQRAYALDTEVAASCGGVVLVGLMALTLSPPYKRYISWCMWWLQYFLTRVEAQ
LHVWVPPLNVRGGRDAVILLMCVVHPTLVFDITKLLLAIFGPLWILQASLLKVPYFVR
VQGLLRICALARKIAGGHYVQMAIIKLGALTGTYYVNHLPTRDWAHNGLRDLAVAVE
PVVFSRMETKLITWGADTAACGDIINGLPVSARRGQEILLGPADGMVSKGWRLLAPIT
AYAQQTRGLLGCIITSLTGRDKNQVEGEVQIVSTATQTFLATCINGVCWTVYHGAGTR
TIASPKGPVIQMYTNVDQDLVGWPAPQGSRSITPCTCGSSDLYLVTRHADVIPVRRRG
DSRGSLLSPRPISYLGSSGGPLLCPAGHAVGLFRAAVCTRGVAKAVDFIPVENLGT
MRSPVFTDNSSPPAVPQSFQVAHLHAPTGSKGSTKVPAAAYAAQGYKVLVLPNSVAATL
GFGAYMSKAHGVDPNIRTGVRTITTGSPITYSTYKFLADGGCSCGGAYDIIICDECHS
TDATSILGIGTVLDQAETAGARLVVLATATPPGSVTVSHPNIEEVALSTTGEIPFYGK
AIPLEVIKGGRHILFCHSKKKCDELAACKLVALGINAVAYYRGLDVSVIPTSGDVVVVS
TDALMTGFTGDFDSVIDCNTCVTQTVDFSLDPTFTIETTTLPQDAVSRTQRRGRTGRG
KPGIYRFVAPGERPSGMFDSSVLCECYDAGCAWYELTPAETTVRLRAYMNTPLPVCQ
DHLEFWEGVFTGLTHIDAHFLSQTKQSGENFPYLVAYQATVCARAQAPPPSWDQMWKC
LIRLKPTLHGPTPLLYRLGAVQNEVTLTHPITKYIMTCMSADLEVVTSTWVLVGGVLA

Fig. 13E

33/38

ALAAYCLSTGCVVIVGRIVLSGKPAIIPDREVLVQEFDEMEEC SQHLPYIEQGMMLAE
QFKQKALGLLQTASRHA EVITPAVQTNWQKLEVFwakHMWNFISGIQYLAGLSTLPGN
PAIASLMAFTAAVTSPLTTGQTLLENILGGWVAAQLAAPGAATAFVGAGLAGAAIGSV
GLGKVLVDILAGYGAGVAGALVAFKIMSGEVPSTEDLVNLLPAILSPGALVVGVCVAA
ILRRHVGPGE GAVQWMNRLIAFASRGNHVSPTHYVPESDAAARVTAISSLTVTQLLR
RLHQWISSECTTPCSGSWLRDIWDWICEVLSDFKTWLKAKLMPQLPGIPFVSCQRGYR
GVWRGDGIMHTRCHCGAEITGHVKNGTMRIVGPRTCRNMWSGTFPINAYTTGPCTPLP
APNYKFALWRVSAEEYVEIRRVGDFHYVSGMTTDNLKPCQIPSPFEFFTELDGVR LHR
FAPPCKPLLREEVSFRVGLHEY PVGSQLPCEPEPDVAVLTSMLTDP SHITAEAAAGRRL
ARGSPPSMASSSASQLSAPSLKATCTANHDS PDAELIEANLLWRQEMGGNITRVESEN
KVVILDSFDPLVAEEDEREVSVP AEILRKSRRFARALPVWARPDYNPPLVETWKKPDY
EPPVVGHCPLPPPRSPVPPPRKKRTVVLTES TLSTALAE LTKSFGSSSTSGITGDN
TTTSSEPAPSGCPPDS DVESYSSMPPLEGE PGDPDLS DGSWSTVSSGADTEDVCCSM
SYSWTGALVTPCAAE EQKL PINALSNSLLRHHNLVYSTTSRSACQRQKKVTFDR LQVL
DSHYQDVLKEVKAAASKVKANLLSVEEACSLTPPHSAKSKFGYGAKDVRCHARKAVAH
INSVWKD LLED SVTPIDTTIMAKNEVF CVQPEKGGRKPARLIVFPDLGVRVCEKMALY
DVVSKLPLAVMGSSYGFQYSPGQRVEFLVQAWKSKKTPMGFSYDTRCFDSTVTESDIR
TEEAIYQCCDLDPQARVAIKSLTERLYVGGPLTNSRGENCYRRCRASGVLT TSCGNT
LTCYIKARAACRAAGLQDCTMLVCGDDL VVICESAGVQEDAASLRAFTEAMTRY SAPP
GDPPQPEYDLELITSCSSNVSVAH DGAGKR VYYLTRDPTT PLARA AWETARHTPVNSW
LGNII MFAPTLWARMILMTHFFSVLIARDQLEQALNCEIYGACYSIEPLDLPPIIQRL
HGLSAFSLHSYSPGEINRVAACLRLKLGVPPLRAWRHRARSVRARLLSRGGRAAICGKY
LFNWAVRTKLKLTPIAAAGRLDLSGWFTAGYSGGDIYHSVSHARPRWFWFCLLLLAAG
VGIYLLPNR"

Fig. 14A

34/38

```

1 gccagccccc tgatgggggc gacactccac catgaatcac tcccctgtga ggaactactg
61 tcttcacgca gaaagcgtct agccatggcg ttagtatgag tgtcgtgcag cctccaggag
121 cccccctccc gggagagcca tagtggtctg cggaaccggg gagtacaccg gaattgccag
181 gacgaccggg tcctttcttg gataaaccgg ctcaatgcct ggagatttgg gcgtgcccc
241 gcaagactgc tagccgagta gtgttgggtc gcgaaaggcc ttgtggtact gcctgatagg
301 gtgcttgcca gtgccccggg aggtctcgta gaccgtgcac catgagcacg aatcctaaac
361 ctcaaagaaa aaccaaacgt aacaccaacc gtcgcccaca ggacgtcaag tcccgggtg
421 gcggtcagat cgttgggtga gtttacttgt tgccgcgcag gggccctaga ttgggtgtgc
481 gcgcgacgag gaagacttcc gagcggtcgc aacctcgagg tagacgtcag cctatcccca
541 aggcacgtcg gcccgagggc aggacctggg ctacgcccgg gtacccttgg cccctctatg
601 gcaatgaggg ttgcgggtgg gcgggatggc tcctgtctcc ccgtggctct cggcctagct
661 ggggccccac agacccccgg cgtaggtcgc gcaatttggg taaggtcatc gataccctta
721 cgtgcggctt cgccgacctc atggggtaca taccgctcgt cggcgccccc cttggaggcg
781 ctgccagggc cctggcgcat ggcgtccggg ttctggaaga cggcgtgaac tatgcaacag
841 ggaaccttcc tggttgctct ttctctatct tccttctggc cctgctctct tgcctgactg
901 tgcccgttcc agcctaccaa gtgcgcaatt cctcggggct ttaccatgtc accaatgatt
961 gccctaactc gagtgttgtg tacgaggcgg ccgatgccat cctgcacact ccggggtgtg
1021 tcccttgctg tcgcgagggt aacgcctcga ggtgttgggt ggcggtgacc cccacggtgg
1081 ccaccaggga cggcaaacct cccacaacgc agcttcgacg tcatatcgat ctgcttgtcg
1141 ggagcgccac cctctgctcg gccctctacg tgggggacct gtgcgggtct gtctttcttg
1201 ttggtcaact gtttaccttc tctcccaggc accactggac gacgcaagac tgcaattggt
1261 ctactatcc cgccatata acgggtctac ccatggcatg gaatatgatg atgaactggg
1321 cccctacggc agcgttgggt gtgctcagc tgctccgaat cccacaagcc atcatggaca
1381 tgatcgctgg cgccactgg ggagtccctg cgggcataaa gtatttctcc atgggtggga
1441 actgggcgaa ggtcctggta gtgctgctgc tatttgccgg cgtcgacgcg gaaaccacg
1501 tcaccggggg aaatgccggc cgcaccacgg ctgggcttgt tggctctcct acaccaggcg
1561 ccaagcagaa catccaactg atcaacacca acggcagttg gcacatcaat agcagggcct
1621 tgaactgcaa tgaaagcctt aacaccgggt ggttagcagg gctcttctat cagcacaat
1681 tcaactcttc aggtgtcct gagaggttgg ccagctgccg acgccttacc gattttgccc
1741 agggctgggg tcctatcagt tatgccacg gaagcggcct cgacgaacgc ccctactgct
1801 ggcactaccc tccaagacct tgtggcattg tgcccgcaa gagcgtgtgt ggcccggat
1861 attgcttcac tcccagcccc gtggtggtgg gaacgaccga caggtcgggc gcgcctacct
1921 acagctgggg tgcaaatgat acggatgtct tcgtccttaa caacaccagg ccaccgctgg
1981 gcaattgggt cggttgatcc tggatgaact caactggatt caccaaagtg tgcggagcgc
2041 ccccttggtg catcgagggt gtgggcaaca acaccttgc ctgccccact gattgcttcc
2101 gcaaatatcc ggaagccaca tactctcggg gcggctccgg tcccaggatt acaccagggt
2161 gcatggtcga ctaccctgat aggttttggc actatccttg taccatcaat tacaccatat
2221 tcaaagtcag gatgtacgtg ggaggggtcg agcacaggct ggaagcggcc tgcaactgga
2281 cgcggggcga acgctgtgat ctggaagaca gggacaggct cgagctcagc ccgttgcctg
2341 tgtccaccac acagtggcag gtccttcctg gttctttcac gacctgcca gccttgtcca
2401 ccggcctcat ccacctccac cagaacattg tggacgtgca gtactgttac ggggtagggt
2461 caagcatcgc gtctgggccc attaagtggg agtacgtcgt tctcctgttc cttctgcttg
2521 cagacgcgcg cgtctgttcc tgcttgggga tgatgttact catatcccaa gcggagggcg
2581 ctttggagaa cctcgtaata ctcaatgcag catccctggc cgggacgcat ggtcttgtgt
2641 ccttctcctg gttcttctgc tttgcgtggg atctgaaggg taggtgggtg cccggagcgg
2701 tctacgccct ctacgggatg tgccctctcc tcctgctcct gctggcgttg gctcagcggg
2761 catacgcact ggacacggag gtggccgcgt cgtgtggcgg cgttgttctt gtcgggttaa
2821 tggcgctgac tctgtcgcca tattacaagc gctatatcag ctggtgcacg tgggtgcttc
2881 agtattttct gaccagagta gaagcgcaac tgcacgtgtg ggttccccc ctcaacgtcc
2941 gggggggggc cgatgccgtc atcttactca cgtgtgtagt acaccgggcc ctgggtattg
3001 acatcaccaa actactcctg gccatcttcg gaccttttg gattcttcaa gccagtttgc
3061 ttaaagtccc ctacttcgtg cgcgttcaag gccttctccg gatctgcgcg ctacgcgcga

```

Fig. 14B

35/38

```

3121 agatagccgg aggtcattac gtgcaaatgg ccatcatcaa gttaggggcg cttactggca
3181 cctgtgtgta taaccatctc gtcctctctc gagactgggc gcacaacggc ctgcgagatc
3241 tggcgtggc tgtggaacca gtcgtctctc cccgaatgga gaccaagctc atcacgtggg
3301 gggcagatac cgccgcgtgc ggtgacatca tcaacggctt gcccgctctc gcccgtaggg
3361 gccaggagat actgcttggg ccagccgacg gaatggtctc caaggggtgg aggttgctgg
3421 cgcccatcac ggcgtacgcc cagcagacga gaggcctcct aggggtgtata atcaccagcc
3481 tgactggccg ggacaaaaac caagtggagg gtgagggtcca gatcgtgtca actgctaccc
3541 agaccttcct ggcaacgtgc atcaatgggg tatgctggac tgtctaccac ggggccggaa
3601 cgaggaccat cgcatacccc aagggtcctg tcatccagac gtataccaat gtggatcaag
3661 acctcgtggg ctggccccgct cctcaagggt cccgctcatt gacacctgac acctgcggtc
3721 cctcggacct ttacctggtc acgaggcacg ccgatgtcat tcccgtgcgc cggcgagggtg
3781 atagcagggg tagcctgctt tcgccccggc ccatttccta ctgaaaggc tcctcggggg
3841 gtccgctgtt gtgccccacg ggacacggcg tgggcctatt cagggccgcg gtgtgcaccc
3901 gtggagtggc taaggcgggtg gactttatcc ctgtggagaa cctagagaca accatgagat
3961 ccccggtgtt cccgacaac tcctctccac cagcagtgcc ccagagcttc caggtggccc
4021 acctgcattg tcccacggc agcggtaaga gcaccaaggc cccggtgcg gaggttgcca
4081 agggctacaa ggtgttgggtg ctcaaccctt ctgttgcctg aacactgggc ttgtgtgctt
4141 acatgtccaa ggcccatggg gttgatccta atatcaggac cggggtgaga acaattacca
4201 ctggcagccc catcacgtac tccacctacg gcaagttcct tgccgacgcc ggggtgctcag
4261 gaggtgctta tgacataata atttgtgacg agtgccactc cacgatgcc acatccatct
4321 cgggcatcgg cactgtcctt gaccaagcag agactgcggg ggcgagactg gttgtgctcg
4381 ccactgctac ccctccgggc tccgtcactg tgtcccatcc taacatcgag gaggttgctc
4441 tgtccaccac cggagagatc cccttttacg gcaaggctat cccctcagag gtgatcaagg
4501 ggggaagaca tctcatcttc tgccactcaa agaagaagtg cgacgagctc gccgcgaagc
4561 tggtcgcatt gggcatcaat gccgtggcct actaccgagg tcttgacgtg tctgtcatcc
4621 cgaccagcgg cgatgttgctc gtcgtgtcga ccgatgctct catgactggc ttaccggcg
4681 acctgcactc tgtgatagac tgcaacacgt gtgtcactca gacagtcatg tttagccttg
4741 acctacctt taccattgag acaaccacgc tccccagga tgctgtctcc aggatcaac
4801 gccggggcag gactggcagg gggaagccag gcatctatag atttggtgca ccgggggagc
4861 gccctccgg catgttcgac tcgtccgtcc tctgtgagtg ctatgacgcg ggctgtgctt
4921 ggtatgagct cacgcccgcg gagactacag ttaggctacg agcgtacatg aacaccccg
4981 ggcttcccg gtgccaggac catcttggtt tttgggaggg cgtctttacg ggcctcactc
5041 atatatagtc ccactttcta tcccagacaa agcagagtgg ggagaacttt ccttacctgg
5101 tagcgtacca agccaccgtg tgcgttaggg ctcaagcccc tcccccatcg tgggaccaga
5161 tgcggaagtg tttgatccgc cttaaaccca ccctccatgg gccaacaccc ctgctataca
5221 gactgggcgc tgttcagaat gaagtcaccc tgacgcaccc aatcaccaaa tacatcatga
5281 catgcatgtc ggccgacctg gaggtcgtca cgagcacctg ggtgctcgtt ggcggcgtcc
5341 tggctgctct ggccgcgtat tgcctgtcaa caggctgctt ggtcatagtg ggcaggatcg
5401 tcttgtccgg gaagccggca attataacct acagggagggt tctctaccag gagttcgatg
5461 agatggaaga gtgctctcag cacttacgtc acatcgagca agggatgatg ctcgctgagc
5521 agttcaagca gaaggccctc ggccctcctg agaccgcgtc ccgccatgca gaggttatca
5581 cccctgctgt ccagaccaac tggcagaaac tcgaggctct ttgggcgaag cacatgtgga
5641 atttcatcag tgggatacaa tacttggcgg gcctgtcaac gctgcctggt aaccccgcc
5701 ttgcttcatt gatggctttt acagctgccg tcaccagccc actaaccact ggccaaaccc
5761 tcctcttcaa catattgggg ggggtgggtg ctgcccagct cgcgcccccc ggtgccccta
5821 ccgcctttgt gggcgctggc ttagctggcg ccgactcga cagcgttgga ctggggaagg
5881 tcctcgtgga cattcttgca ggcctatggc cggcgctggc gggagctctt gtggcattca
5941 agatcatgag cggtgaggct ccctccacgg aggcctggt caatctgctg cccgccatcc
6001 tctcacctgg agcccttgca gtcggtgtgg tctttgcatc aatactgcgc cggcgtgttg
6061 gccggggcga gggggcagtg caatggatga accggcta at agccttcgcc tcccggggga
6121 accatgtttc cccacacac tacgtgccgg agagcgatgc agccgcccgc gtcactgcca
6181 tactcagcag cctcactgta acccagctc tgaggcgact gcatcagtgg ataagctcg
6241 agtgataccac tccatgctcc ggttcctggc taagggacat ctgggactgg atatgcgagg
6301 tgctgagcga ctttaagacc tggctgaaag ccaagctcat gccacaactg cctgggattc
6361 cctttgtgtc ctgccagcgc gggatatagg gggctgtggc aggagacggc attatgcaca
6421 ctcgtgcca ctgtggagct gagatcactg gacatgtcaa aaacgggacg atgaggatcg
6481 tcggtcctag gacctgaag aacatgtgga gtgggacgtt cttcattaat gcctacacca

```

Fig. 14C

36/38

6541	cgggccctg	tactccctt	cctgcgccga	actataagtt	cgcgctgtg	aggggtgtctg
6601	cagaggaata	cgtggagata	aggcgggtg	gggacttcca	ctacgtatcg	ggcatgacta
6661	ctgacaatct	caaatgccc	tgccagatcc	catcgcccga	atttttcaca	gaattggacg
6721	gggtgcgcct	acataggttt	gcgccccctt	gcaagccctt	gctgcgggag	gaggtatcat
6781	tcagagtagg	actccacgag	tacccggttg	ggtcgcaatt	accttgcgag	cccgaaccgg
6841	acgtagccgt	gttgacgtcc	atgctcactg	atccctccca	tataacagca	gaggcggccg
6901	ggagaaggtt	ggcgagaggg	tcacccccctt	ctatggccag	ctcctcggct	agccagctgt
6961	ccgctccatc	tctcaaggca	acttgacccg	ccaacatga	ctcccctgac	gccgagctca
7021	tagaggctaa	cctcctgtgg	aggcaggaga	tgggcgga	catcaccagg	gttgagtcag
7081	agaacaaagt	ggtgattctg	gactccttcg	atccgcttgt	ggcagaggag	gatgagcggg
7141	aggtctccgt	acccgcagaa	attctgcgga	agtctcgag	attcgcccc	gccctgccc
7201	tctgggcgcg	gccggactac	aacccccctg	tagtagagac	gtggaaaaag	cctgactacg
7261	aaccacctgt	ggtccatggc	tgcccgctac	cacctccacg	gtccccctct	gtgcctccgc
7321	ctcgaaaaa	gcgtacggtg	gtcctcaccg	aatcaaccct	acctactgcc	ttggccgagc
7381	ttgccaccaa	aagttttggc	agctcctcaa	cttcggcat	tacgggcgac	aatacgacaa
7441	catcctctga	gcccccccc	tctggctgcc	ccccgactc	cgacgttgag	tcctattctt
7501	ccatgcccc	cctggagggg	gagcctgggg	atccggatct	cagcgacggg	tcatggctga
7561	cggtcagtag	tggggccgac	acggaagatg	tcgtgtgctg	ctcaatgtct	tattcctgga
7621	caggcgcact	cgtcaccccc	tgcgtgcgg	aggaacaaaa	actgcccac	aacgcactga
7681	gcaactcgtt	gtacgccat	cacaatctgg	tgtattccac	cacttcacgc	agtgtctg
7741	aaaggaagaa	gaaagtcaca	tttgacagac	tgcaagttct	ggacagccat	taccaggacg
7801	tgctcaagga	ggtcaaagca	gcggcgtaaa	aagtgaaggc	taacttgcta	tccttagagg
7861	aagcttgag	cctggcgccc	ccacattcag	ccaaatccaa	gtttggctat	ggggcaaaaag
7921	acgtccgttg	ccatgccaga	aaggccgtag	cccacatcaa	ctccgtgtgg	aaagaccttc
7981	tgaagacag	tgtaacacca	atagacacta	ccatcatggc	caagaacgag	gttttctg
8041	ttcagcctga	gaaggggggt	cgtaagccag	ctcgtctcat	cgtgttcccc	gacctggg
8101	tgcgcgtgtg	cgagaagatg	gccctgtacg	acgtgggttag	caagctcccc	ttggccgtga
8161	tgggaagctc	ctacggattc	caatactcac	caggacagcg	ggttgaattc	ctcgtgcaag
8221	cgtggaagtc	caagaagacc	ccgatggggc	tctcgtatga	taccgctgt	tttgactcca
8281	cagtcactga	gagcgacatc	cgtacggagg	aggcaattta	ccaatgttgt	gacctggacc
8341	cccaagcccc	cgtggccatc	aagtcctca	ctgagaggct	ttatgttggg	ggccctctta
8401	ctaattcaag	gggggaaaac	tgccggtacc	gcaggtgccg	cgcgagcaga	gtactgacaa
8461	ctagctgtgg	taacaccctc	actcgtaca	tcaaggccc	ggcagcctgt	cgagccgcag
8521	ggctccagga	ctgcaccatg	ctcgtgtgtg	gcgacgaact	agtcgttatc	tgtgaaagt
8581	cggggggtcca	ggaggacgcg	gcgagcctga	gagccttcac	ggaggctatg	accagggtact
8641	ccgccccccc	cggggacccc	ccacaaccag	aatacgactt	ggagcttata	acatcatgct
8701	cctccaacgt	gtcagtcgcc	cacgacggcg	ctggaaagag	ggtctactac	cttaccctgt
8761	accctacaac	ccccctcgcg	agagccgcgt	gggagacagc	aagacacact	ccagtcaatt
8821	cctggctagg	caacataatc	atgtttgccc	ccacactgtg	ggcgaggatg	atactgatga
8881	ccacttctt	tagcgtcctc	atagccagg	atcagcttga	acaggctctc	aactgcgaga
8941	tctacggagc	ctgctactcc	atagaaccac	tggatctacc	tccaatcatt	caaagactcc
9001	atggcctcag	cgcattttca	ctccacagtt	actctccagg	tgaatataat	aggggtggccg
9061	catgcctcag	aaaacttggg	gtcccgccct	tgcgagcttg	gagacaccgg	gcctggagcg
9121	tccgcgctag	gcttctggcc	agaggaggca	aggctgcca	atgtggcaag	tacctcttca
9181	actgggcagt	aagaacaaa	ctcaaaactca	ctccgataac	ggccgctggc	cggctggact
9241	tgtccgctg	gttcacggct	ggctacagcg	ggggagacat	ttatcacagc	gtgtctcatg
9301	cccgcccccg	ctggttctgg	ttttgcctac	tctgtcttgc	tgcaggggta	ggcatctacc
9361	tcctcccaaa	ccgatgaaga	ttgggctaac	cactccaggc	caataggcca	ttccct

Fig. 14D

37/38

MSTNPKPQRKTKRNTNRRPQDVKFPGGGQIVGGVYLLPRRGPRL
GVRATRKTSESRQPRGRRQPIPKARRPEGRTWAQPGYPWPLYGNEGCGWAGWLLSPRG
SRPSWGPTDPRRRSRNLGKVIDTLTCGFADLMGYIPLVGAPLGGAARALAHGVRVLED
GVNYATGNLPGCSFSIFLLALLSCLTVPASAYQVRNSSGLYHVTNDCPNSSVVEAAD
AILHTPGCVPCVREGNASRCWVAVTPTVATRDGKLPTTQLRRHIDLLVGSATLCSALY
VGDLGGSVFLVGQLFTFSPRHHWTTQDCNCSIYPGHITGHRMAWNMMMNWSPTAALVV
AQLLRIPQAIMDMIAGAHWGVLAGIKYFSMVGWNAKVLVVLFFAGVDAETHVTGGNA
GRTTAGLVGLLTPGAKQNIQLINTNGSWHINSTALNCNESLNTGWLAGLFYQHKFNSS
GCPERLASCRRLTDFAGWGPISYANGSGLDERPYCWHYPPRPGIVPAKSVCGPVYC
FTSPVVVGTTDRSGAPTYSWGANDTDVFLNNTRPPLGNWFGCTWMNSTGFTKVCGA
PPCVIGGVGNNTLLCPTDCFRKYPEATYSRCGSGPRITPRCMVDYPYRLWHYPCTINY
TIFKVRMYVGGVEHRLEAACNWTGRGERCDLEDRDRSELSPLLLSTTQWQVLPCSFTTL
PALSTGLIHLHQNIVDVQYLYGVGSSIASWAIKWEYVVLFLLLADARVCSCWMLL
ISQAEAALENLVILNAASLAGTHGLVSFLVFCCFAWYKGRWVPGAVYALYGMWPLLL
LLLALPQRAYALDEVAASCGGVVLVGLMALTSPYKRYISWCMWWLQYFLTRVEAQ
LHVWVPPLNVRGGRDAVILLTCVHPALVFDITKLLLAIFGPLWILQASLLKVYPFVR
VQGLLRICALARKIAGGHYVQMAIIKLGALTGTCVYNHLAPLRDWAHNGLRDLAVAVE
PVVFSRMETKLITWGADTAACGDIINGLPVSARRGQEILLGPADGMVSKGWRLAPIT
AYAQQTRGLLGCIITSITGRDKNQVEGEVQIVSTATQTFLATCINGVCWTVYHGAGTR
TIASPKGPVIQTYTNVDQDLVGWPAPQGSRLTPCTCGSSDLYLVTRHADVIPVRRRG
DSRGSLLSPRPISYKGS GG PLLCPTGHAVGLFRAAVCTRGVAKAVDFIPVENLETT
MRSPVFTDNSSPPAVPQSFQVAHLHAPTSGSKSTKVPAAAYAAKGYKVLVLNPSVAATL
GFGAYMSKAHGVDPNIRTGVRTITTGSPITYSTYGKFLADAGCSGGAYDIIICDECHS
TDATSIISGIGTVLDQAETAGARLVVLATATPPGSVTVSHPNIEEVALSTTGEIPFYGK
AIPLEVIKGRHLIFCHSKKKCDELAALVALGINAVAYYRGLDVSVIPTSGD VVVVS
TDALMTGFTGDFDSVIDCNTCVTQTVD FSLDPTFTIETTTLPQDAVSRTQRRGRTGRG
KPGIYRFVAPGERPSGMFDSSVLCECYDAGCAWYELTPAETTVRLRAYMNT PGLPVCQ
DHLGFWEGVFTGLTHIDAHFLSQTKQSGENFPYLVAYQATVCARAQAPPPSWDQMRKC
LIRLKPTLHGPTPLLYRLGAVQNEVTLTHPITKYIMTCMSADLEVVTSTWVLVGGVLA

Fig. 14E

38/38

ALAAYCLSTGCVVIVGRIVLSGKPAIIPDREVLVQEFDEMEEC SQHLPYIEQGMLAE
QFKQKALGLLQTASRHAEVITPAVQTNWQKLEFWAKHMWNFISGIQYLAGLSTLPGN
PAIASLMAFTAAVTSPLTTGQTLLFNILGGWVAAQLAAPGAATAFVGAGLAGAALDSV
GLGKVLVDILAGYGAGVAGALVAFKIMSGEVPSTEDLVNLLPAILSPGALAVGVVVFAS
ILRRRVGPGEGAVQWMNRLIAFASRGNHVSPTHYVPESDAAARVTAILSSLTQLLR
RLHQWISSECTTPCSGSWLRDIWDWICEVLSDFKTWLKAKLMPQLPGIPFVSCQRGYR
GVWRGDGIMHTRCHCGAEITGHVKNGTMRIVGPRTCKNMWSGTTFFINAYTTGPCTPLP
APNYKFALWRVSAEEYVEIRRVGDFHYVSGMTTDLNLCPCQIPSPFEFFTEL DGVRLHR
FAPPCKPLLREEVSFRVGLHEYVPGSQLPCEPEPDVAVLTSM L TDP SHITAE AAGRRL
ARGSPPSMASSSASQLSAPSLKATCTANHDS PDAELIEANLLWRQEMGGNITRVESEN
KVVILDSFDPLVAEEDEREVSVP AEILRKSRRFAPALPVWARPDYNPLL VETWKKPDY
EPPVVHGCPLPPPRSPVP PPRKRTVVLTESTLPTALAE L ATKSF GSSSTSGITGDN
TTTSSEPAPSGCPPDSDVESYSSMPLEGE PGDPDLS DGSWSTVSSGADTEDVCCSM
SYSWTGALVTPCAAEEQKL PINALSNSLLRHHNLVYSTTSRSACQRKKKVT FDR LQVL
DSHYQDVLKEVKAAASKVKANLLSVEEACSLAPPHSAKSKFGYGAKDVRCHARKAVAH
INSVWKDLLED SVTPIDTTIMAKNEVFCVQPEKGGRKPARLIVFPDLGVRVCEKMALY
DVVSKLPLAVMGSSYG FQYSPGQ RVEFLVQAWKSKKTPMGLSYDTRCFDSTVTESDIR
TEEAIYQCCDLDPQARVAIKSLTERLYVGGPLTNSRGENCGYRRCRASRVLT TSCGNT
LTRYIKARAACRAAGLQDCTMLVCGDDL V VICESAGVQEDAASLRAFTEAMTRY SAPP
GDPPQPEYDLELITSCSSNVSV AHDGAGKRVYYLTRDPTT PLARA AWETARHTPVNSW
LGN IIMFAPTLWARMILMTHFFSVLIARDQLEQALNCEIYGACYSIEPLDLPP I IQRL
HGLSAFSLHSYSPGEINRVAACLRLGVPPLRAWRHRAW SVRARLLARGGKAAICGKY
LFNWAVRTKLKLT PITAAGRLDL SGWFTAGYSGGDIYHSVSHARPRWFWFCLLLLAAG
VGIYLLPNR"

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
16 June 2005 (16.06.2005)

PCT

(10) International Publication Number
WO 2005/053516 A3

(51) International Patent Classification⁷: **C07H 21/02**,
21/04, C12P 19/34, 21/00, C12N 15/09, 15/82, 15/85,
15/00, C12Q 1/70, 1/68

(21) International Application Number:
PCT/US2004/040120

(22) International Filing Date: 1 December 2004 (01.12.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/525,989 1 December 2003 (01.12.2003) US

(71) Applicant (*for all designated States except US*): **BOARDS OF REGENTS, THE UNIVERSITY OF TEXAS SYSTEM** [US/US]; 201 West 7th Street, Austin, TX 78701 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **LEMON, Stanley, M.** [US/US]; 1517 Bayou Shore Drive, Galveston, TX 77551 (US). **YI, MinKyung** [KR/US]; 7700 Seawall Blvd.#301, Galveston, TX 77551 (US).

(74) Agent: **PROVENCE, David, L.**; Mueting, Raasch & Gebhardt, P.A., P.O. Box 581415, Minneapolis, MN 55454-1415 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
29 December 2005

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: REPLICATION COMPETENT HEPATITIS C VIRUS AND METHODS OF USE

(57) Abstract: The present invention provides replication competent polynucleotides that include a coding sequence encoding a hepatitis C virus polyprotein having adaptive mutations. The invention also includes methods for making replication competent polynucleotides, identifying a compound that inhibits replication of a replication competent polynucleotide, selecting a replication competent polynucleotide, and detecting a replication competent polynucleotide.



WO 2005/053516 A3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US04/40120

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07H 21/02, 21/04; C12P 19/34, 21/00; C12N 15/09, 15/82, 15/85, 15/00; C12Q 1/70, 1/68

US CL : 536/23.1, 23.72; 435440, 441, 442455, 91.1, 91.4, 91.4291.51, 69.1, 69.270.1, 455, 5, 6, 94

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1, 23.72; 435440, 441, 442455, 91.1, 91.4, 91.4291.51, 69.1, 69.270.1, 455, 5, 6, 94

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	US6,689,559B2 (WIMMER et al) 10 February 2004 (10.02.2004), Fig. 2-4, Claims 1-12, columns 11-13).	1-36
X	US 20020098202A1 (WIMMER et al) 24 October 2002 (24.10.2002), see Fig. 1-4, claims 1-52)	1-36
A	IKEDA, M. et al. Selection Subgenomic and Genome-Length Dicistronic RNAs Derived from an Infectious Molecular Clone of the HCV-N Strain of hepatitis C virus Replicate Efficiently in Cultured Huh7 Cells. J. Virol. March 2002, Vol. 76. No. 6, pages 2997-3006.	
A	LOHMANN, V. et al. Mutations in Hepatitis C Virus RNAs conferring Cell Culture Adaptation. J. Virol. February 2001, Vol. 75. No. 3, pages 1437-1449.	
X	BUKH, J. et al. Mutations that permit efficient replication of hepatitis C virus RNA in Huh-7 cells prevent productive replication in chimpanzees, PNAS, October 2002, Vol. 99. No. 22, pages 14416-14421, see page 14416, 2nd column, last paragraph and 1st column, 2nd paragraph.	1-22, 7, 9-10, 14-17,
A	KRIEGER, N. et al. Enhancement of Hepatitis C Virus RNA Replication by Cell Culture-Adaptive Mutations. J. Virol. May 2001, Vol. 75. No. 10, pages 4614-4624.	

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

Special categories of cited documents:

"A"	document defining the general state of the art which is not considered to be of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier application or patent published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed	"&"	document member of the same patent family

Date of the actual completion of the international search

17 August 2005 (17.08.2005)

Date of mailing of the international search report

21 SEP 2005

Name and mailing address of the ISA/US

Mall Stop PCT, Attn: ISA/US

Commissioner for Patents

P.O. Box 1450

Alexandria, Virginia 22313-1450

Facsimile No. (703) 305-3230

Authorized officer

Bao Qun Li

Telephone No. 571-272-1600

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US04/40120

Continuation of B. FIELDS SEARCHED Item 3:
WEST, MEDLINE, CAPLUS
Search terms: HCV replicon, in vitro, synthesis, construct, expression system